

# Effect of Nebulized Magnesium vs Placebo Added to Albuterol on Hospitalization Among Children With Refractory Acute Asthma Treated in the Emergency Department

## A Randomized Clinical Trial

Suzanne Schuh, MD; Judy Sweeney, RN, BScN; Maggie Rumantir, MD; Allan L. Coates, MDCM, BEng; Andrew R. Willan, PhD; Derek Stephens, MSc, BSc; Eshetu G. Atenafu, MSc; Yaron Finkelstein, MD; Graham Thompson, MD; Roger Zemek, MD; Amy C. Plint, MD, MSc; Jocelyn Gravel, MD, MSc; Francine M. Ducharme, MD, MSc; David W. Johnson, MD; Karen Black, MD, MSc; Sarah Curtis, MD; Darcy Beer, MD; Terry P. Klassen, MD, MSc; Darcy Nicky, BSc, PhM; Stephen B. Freedman, MDCM, MSc; for the Pediatric Emergency Research Canada (PERC) Network

**IMPORTANCE** While intravenous magnesium decreases hospitalizations in refractory pediatric acute asthma, it is variably used because of invasiveness and safety concerns. The benefit of nebulized magnesium to prevent hospitalization is unknown.

**OBJECTIVE** To evaluate the effectiveness of nebulized magnesium in children with acute asthma remaining in moderate or severe respiratory distress after initial therapy.

**DESIGN, SETTING, AND PARTICIPANTS** A randomized double-blind parallel-group clinical trial from September 26, 2011, to November 19, 2019, in 7 tertiary-care pediatric emergency departments in Canada. The participants were otherwise healthy children aged 2 to 17 years with moderate to severe asthma defined by a Pediatric Respiratory Assessment Measure (PRAM) score of 5 or greater (on a 12-point scale) after a 1-hour treatment with an oral corticosteroid and 3 inhaled albuterol and ipratropium treatments. Of 5846 screened patients, 4332 were excluded for criteria, 273 declined participation, 423 otherwise excluded, 818 randomized, and 816 analyzed.

**INTERVENTIONS** Participants were randomized to 3 nebulized albuterol treatments with either magnesium sulfate (n = 410) or 5.5% saline placebo (n = 408).

**MAIN OUTCOMES AND MEASURES** The primary outcome was hospitalization for asthma within 24 hours. Secondary outcomes included PRAM score; respiratory rate; oxygen saturation at 60, 120, 180, and 240 minutes; blood pressure at 20, 40, 60, 120, 180, and 240 minutes; and albuterol treatments within 240 minutes.

**RESULTS** Among 818 randomized patients (median age, 5 years; 63% males), 816 completed the trial (409 received magnesium; 407, placebo). A total of 178 of the 409 children who received magnesium (43.5%) were hospitalized vs 194 of the 407 who received placebo (47.7%) (difference, -4.2%; absolute risk difference 95% [exact] CI, -11% to 2.8%);  $P = .26$ . There were no significant between-group differences in changes from baseline to 240 minutes in PRAM score (difference of changes, 0.14 points [95% CI, -0.23 to 0.50];  $P = .46$ ); respiratory rate (0.17 breaths/min [95% CI, -1.32 to 1.67];  $P = .82$ ); oxygen saturation (-0.04% [95% CI, -0.53% to 0.46%];  $P = .88$ ); systolic blood pressure (0.78 mm Hg [95% CI, -1.48 to 3.03];  $P = .50$ ); or mean number of additional albuterol treatments (magnesium: 1.49, placebo: 1.59; risk ratio, 0.94 [95% CI, 0.79 to 1.11];  $P = .47$ ). Nausea/vomiting or sore throat/nose occurred in 17 of the 409 children who received magnesium (4%) and 5 of the 407 who received placebo (1%).

**CONCLUSIONS AND RELEVANCE** Among children with refractory acute asthma in the emergency department, nebulized magnesium with albuterol, compared with placebo with albuterol, did not significantly decrease the hospitalization rate for asthma within 24 hours. The findings do not support use of nebulized magnesium with albuterol among children with refractory acute asthma.

**TRIAL REGISTRATION** ClinicalTrials.gov Identifier: [NCT01429415](https://clinicaltrials.gov/ct2/show/study/NCT01429415)

JAMA. 2020;324(20):2038-2047. doi:10.1001/jama.2020.19839

- [+ Visual Abstract](#)
- [+ Supplemental content](#)
- [+ CME Quiz at \[jamacmelookup.com\]\(http://jamacmelookup.com\)](#)

**Author Affiliations:** Author affiliations are listed at the end of this article.

**Group Information:** The Pediatric Emergency Research Canada (PERC) Network members are listed at the end of the article.

**Corresponding Author:** Suzanne Schuh, MD, Pediatric Emergency Medicine, The Hospital for Sick Children, 555 University Ave, Toronto, M5G 1X8, ON, Canada ([suzanne.schuh@sickkids.ca](mailto:suzanne.schuh@sickkids.ca)).

Acute asthma is a leading cause of pediatric hospitalizations.<sup>1</sup> While inhaled  $\beta_2$  agonists, anticholinergics, and systemic corticosteroids are widely recommended to reduce hospitalizations in severe asthma exacerbations,<sup>2-4</sup> some children have a limited response to  $\beta_2$  agonists and corticosteroids.<sup>5</sup>

When administered intravenously (IV) to children with severe acute asthma, magnesium has been shown to decrease hospitalizations from the emergency department (ED).<sup>6-9</sup> However, children are rarely discharged home after receiving IV magnesium, and IV magnesium is often given with the intent to prevent intensive care admission.<sup>10,11</sup> Clinician reluctance to use IV magnesium may be in part because of the pain associated with IV insertion and because of its well-known association with hypotension.<sup>10,11</sup> In contrast, administration of magnesium by nebulization is noninvasive, and potentially more efficient, because this treatment mode targets the lower airway and reduces systemic exposure.

Previously published pediatric asthma studies of nebulized magnesium were small,<sup>12,13</sup> and had methodological limitations such as lack of focus on health care utilization outcomes<sup>13-15</sup> and not limiting participants to those not responsive to initial optimized care.<sup>12,13,15</sup> Therefore, the Magnesium Nebulization Utilization Management in Pediatric Asthma (MAGNUM PA) Trial was designed to evaluate the effectiveness of inhaled magnesium in children who presented to EDs with an acute asthma exacerbation and remained in moderate or severe respiratory distress after evidence-based standardized initial therapy. The study hypothesis was that the administration of 3 magnesium sulfate nebulization treatments added to ongoing albuterol treatment would result in a lower 24-hour hospitalization rate compared with albuterol nebulization with placebo.

## Methods

### Design

This was a randomized, double-blind, parallel-group, placebo-controlled trial comparing the effect of nebulized magnesium sulfate with albuterol vs placebo with albuterol in children with refractory acute asthma. Written informed consent was obtained from all caregivers and participants, along with assent, as appropriate. The research ethics boards of all institutions approved the trial. The trial was registered at ClinicalTrials.gov and the protocol has been published<sup>16</sup> (and is available in [Supplement 1](#)) and the statistical analysis plan and its amendments are summarized in [Supplement 2](#), [Supplement 3](#), and [Supplement 4](#). In this trial, we enrolled children with acute asthma in 7 Canadian tertiary-care pediatric EDs belonging to the Pediatric Emergency Research Canada Network (eTable 1 in [Supplement 5](#)).

### Trial Participants

Children 2 to 17 years of age were eligible if they had a diagnosis of asthma made by a physician, had a previous episode of acute wheeze treated with an inhaled bronchodilator or a systemic corticosteroid, and had persistent moderate to

## Key Points

**Question** What is the effectiveness of nebulized magnesium in children and adolescents with acute asthma in the emergency department who remain in moderate or severe respiratory distress after evidence-based standardized initial therapy?

**Findings** In this randomized clinical trial that included 816 patients, nebulized magnesium with albuterol, compared with placebo with albuterol, did not significantly decrease the rate of hospitalization for asthma within 24 hours (43.5% vs 47.7%, respectively).

**Meaning** The findings do not support use of nebulized magnesium with albuterol among children with refractory acute asthma.

severe asthma after the completion of a 1-hour initial asthma treatment period. The latter was designed to optimize routine management and included an oral corticosteroid (ie, dexamethasone, 0.3-0.6 mg/kg/dose, maximum, 20 mg; or prednisolone, 1-2 mg/kg/dose, maximum, 60 mg) with 3 consecutive inhaled albuterol (500-1000  $\mu$ g/treatment) and ipratropium bromide (80  $\mu$ g/treatment) treatments via a metered dose inhaler or a nebulizer (albuterol, 5 mg/treatment; ipratropium bromide, 500  $\mu$ g/treatment). We excluded children requiring immediate airway management; children who received IV magnesium prior to enrollment; children with comorbidities such as chronic lung, cardiovascular, kidney, neurologic, or other systemic disease; and children with a known hypersensitivity to magnesium. Families without adequate command of the English or French language, without telephone or e-mail contact information, and those previously enrolled were also excluded.

Persistent moderate to severe asthma after the aforementioned initial therapy was defined by a Pediatric Respiratory Assessment Measure (PRAM) score of 5 points or greater ([Table 1](#); eAppendix 2 in [Supplement 3](#)).<sup>19</sup> This cutoff was chosen based on the high proportion of asthma hospitalizations with PRAM scores of 5 or greater after initial therapy (eTables 2 and 3 in [Supplement 5](#)). The PRAM score represents a 12-point pediatric acute asthma measurement tool (higher PRAM scores indicate higher asthma severity), which has been fully validated in the ED setting for use in children 12 months of age or older.<sup>17,20,21</sup>

### Randomization and Blinding

Using random number-generating software at <https://www.randomizer.org/>, the research pharmacist (D.N.) at the coordinating center produced master randomization tables, stratified by site and age ( $\leq 5$  years vs  $\geq 6$  years). Permuted block randomization with block sizes of 6 and 8 with a 1:1 allocation ratio of magnesium to placebo was used. The randomization sequence was restricted to the coordinating research pharmacy until the study database was locked. Consecutively numbered kits containing either magnesium sulfate or 5.5% saline placebo (to match tonicity of magnesium sulfate) were prepared by each pharmacy according to the procedure manual supplied by the coordinating center.

Table 1. Pediatric Respiratory Assessment Measure Score<sup>a</sup>

Sign	Points			
	0	1	2	3
Suprasternal retractions	Absent		Present	
Scalene muscle contraction	Absent		Present	
Air entry	Normal	Decreased at bases	Widespread decrease	Absent/minimal
Wheezing	Absent	Expiratory only	Inspiratory and expiratory	Audible without stethoscope/silent chest with minimal air entry
Oxygen saturation	≥95%	92%-94%	≤91%	
Oxygen saturation in Calgary <sup>b</sup>	≥93%	90%-92%	≤89%	

Reprinted from *The Journal of Pediatrics* with permission from Elsevier.<sup>17</sup>

<sup>a</sup> PRAM score represents the sum of the individual components (range, 0-12). PRAM scores of 0-3, 4-7, and 8-12 points represent mild, moderate, and severe asthma, respectively.<sup>17</sup>

<sup>b</sup> Because Calgary, Alberta, Canada, is 1000 m above sea level, the saturation cutoffs have been adjusted accordingly<sup>18</sup> (see eAppendix 2 in Supplement 5 for details).

The magnesium and placebo solutions were identical in volume, color, taste, and smell, both in the steady state and during nebulization, as verified by the coordinating center pharmacy. Study participants, research nurses, ED staff, and the study analyst were blinded to the treatment assignment (see eAppendixes 3 and 4 and eTable 4 in Supplement 5 for study blinding and unblinding procedures).

### Procedures

Prior to the study start, dedicated research nurses received training about the study protocol, data collection, and treatment delivery using standardized in-person sessions. They collected patient-level data and study-related outcomes, conducted trial interventions, and entered study data into a secure electronic database. To maximize the accuracy of the PRAM measurement, all study nurses and site investigators completed an online PRAM training module<sup>22</sup> available at <https://enseignement.chusj.org/PRAM-En>.

Eligible participants received 3 consecutive nebulization treatments, consisting of 5 mg (1 mL) of albuterol and either 600 mg (1.2 mL) of magnesium sulfate (Sandoz) or 1.2 mL of 5.5% saline. Sterile water (3.8 mL) was added to each nebulization treatment in both groups to achieve identical tonicity of both intervention and placebo solutions (eAppendix 3 and eTable 4 in Supplement 5). The selected dose of magnesium was at the upper end of the dosage range previously used.<sup>23</sup> Because pulmonary deposition, in terms of dose/kg, of nebulization treatments in younger children is similar to that in their older counterparts,<sup>24</sup> we used the same magnesium dose in all participants. To optimize pulmonary deposition in all participants, we pretested and selected the AeroNebGo nebulizer (Philips) attached to the Idehaler holding chamber (La Diffusion Technique Française). This high-efficiency delivery system has been shown to deliver 20% of the medication dose to the lungs compared with 4% delivered via conventional nebulizers (eAppendix 5 in Supplement 5).<sup>25,26</sup>

Emergency physicians used their clinical judgment to determine the need for additional asthma therapies and made disposition determinations as per usual practice. Discharged children received prescriptions for inhaled albuterol, oral corticosteroid, and inhaled corticosteroid.

### Outcomes

The primary outcome measure was physicians' decision to hospitalize children for persistent respiratory distress or the need for supplemental oxygen within 24 hours of randomization. Hospitalization is a clinically powerful and policy-relevant marker of treatment failure of importance to clinicians, families, and health care organizations because half of pediatric asthma costs relate to hospitalizations.<sup>27</sup>

The secondary outcomes specified a priori included changes in the PRAM score (PRAM is a validated summative pediatric acute asthma score ranging 0-12 points, where PRAM scores of 1-3, 4-7, and 8-12 points indicate mild, moderate, and severe asthma, respectively, with a change of 3 points regarded as clinically significant; Table 1)<sup>17,20</sup>; respiratory rate; oxygen saturation (on room air) from postrandomization baseline to 60, 120, 180, and 240 minutes (or up to the time of disposition) after starting the intervention; changes in blood pressure from baseline to 20, 40, 60, 120, 180, and 240 minutes; and the number of additional albuterol treatments administered within 240 minutes. Adverse effects included adverse events (AEs) coded using the Medical Dictionary for Regulatory Activities, version 19, and serious AEs (SAEs). Predefined expected life/disease-related occurrences, such as cough, respiratory distress, asthma-related hospitalization, IV insertion, sinus tachycardia, and bitter/salty taste of the experimental solution, were not considered AEs in either group and were not measured as such. We a priori defined SAEs as hypotension requiring medical intervention, apnea, or admission to the intensive care unit. Other (exploratory) outcomes included asthma-related (1) hospitalizations, (2) unscheduled medical visits to any facility within 72 hours of ED discharge, and (3) administration of IV magnesium in the ED after the experimental intervention.

### Statistical Analyses

This trial was initially launched as a 2-center trial, with a targeted sample size of 284 patients, which was able to detect a minimally significant difference of 15 percentage points to decrease the hospitalization rate (ie, primary outcome) from 30% to 15%, with a power of 80%. This difference was determined after discussion with all study investigators. However, during this phase, the primary outcome rate had an overall event

rate of 50% (no between-group analysis was performed) and we thus realized we would be underpowered to evaluate our primary outcome. Therefore, this phase of the study was considered to be a pilot phase and the knowledge gained informed the full study design and the final protocol sample size calculations targeting a difference of 10 percentage points between groups in the primary outcome. Because the study remained blinded, and no analyses were performed, the final significance threshold remained unchanged.

The new targeted difference of 10 percentage points was based on a national survey of pediatric emergency medicine physicians<sup>10</sup> and on the evidence that this difference has previously led to changes in national guidelines.<sup>28</sup> Using a type I, 2-sided error of .05 and 80% power, our new targeted sample size was 816 participants.

All analyses were specified a priori and performed according to the assigned randomization group. All significance thresholds were 2-sided. To account for the multiple comparisons required to assess secondary outcomes, we used the Holm method, resulting in a significance level of 0.008 for each of the 6 secondary outcomes.<sup>29</sup> Other outcomes were not adjusted for multiplicity. Because adverse effects were uncommon, these were reported only in a descriptive way.

The data monitoring committee met annually (eAppendix 6 in Supplement 5). There was 1 planned interim analysis after the first 200 randomized patients, with a 1-sided hypothesis for the primary outcome set at 0.01 significance level. Because this 1-tailed test would only reject the null hypothesis if hospitalizations were more frequent in the magnesium than in the placebo group (contrary to the study hypothesis), the significance level for the final analysis of efficacy was not adjusted.

Baseline variables were summarized with descriptive statistics. For our primary analysis, we used a 2-sided Fisher exact test to examine the treatment effect on hospitalization within 24 hours. Further, we performed post hoc analyses using generalized linear mixed modeling for the primary and all other outcome analyses to control for randomization stratification by age group and site, where the site was treated as a random effect. This method was also used in the per-protocol analysis to estimate treatment effect of magnesium in children who received all experimental treatments. Adjusted relative risk differences were used to quantify effect sizes. We used the following a priori-identified subgroups for subgroup analyses: postrandomization baseline PRAM score of 8 or greater (indicating severe asthma),<sup>17,20</sup> age 5 years or younger,<sup>30</sup> male sex,<sup>31</sup> personal history of atopy, and historical report of viral-induced preschool wheeze (defined as age ≤5 years without atopy or cough between upper respiratory infections).<sup>32</sup> For subgroup analyses, we used generalized linear mixed modeling with treatment group-subgroup interaction factor, controlling for stratification variables, and reported adjusted risk differences for each subgroup. To analyze our secondary outcomes, we used the mixed-model method to compare changes from baseline in PRAM score, respiratory rate, oxygen saturation, and blood pressure between groups for the times previously specified, and generalized linear mixed modeling with negative binomial distri-

bution to compare the number of additional albuterol ED treatments administered within 240 minutes between groups. The aforementioned generalized linear mixed modeling was also used to examine magnesium treatment effect on hospitalizations within 72 hours, revisits within 72 hours, and IV magnesium treatment after the experimental therapy.

No imputation was conducted for the missing data. Overall significance for primary and secondary outcomes was set at .05 (2-sided). Statistical analysis was performed using version 9.4 of the SAS system for Windows (SAS Institute, 2002-2012) and the open source statistical software R version 3.5.3 (The R Foundation for Statistical Computing, Vienna, Austria, 2019).

## Results

Between September 26, 2011, and November 19, 2019, 818 participants were randomized. A total of 99.8% (409/410) of children in the magnesium sulfate and 99.8% (407/408) of those in the placebo groups completed follow-up (Figure 1). Because 2 participants could not be included in the primary analysis, our data monitoring committee agreed with expanding enrollment to 818 patients. Trial groups were similar with respect to baseline characteristics (Table 2). Overall, 756 children (93%) received all 3 intervention treatments; 88.3% (361/409) in the magnesium and 97.1% (395/407) in the placebo groups.

Missing data were negligible (<5%) to none, with the exception of the preschool wheeze subgroup (missing data, 12%). Nonimputed data were used in the presented results.

### Primary Outcome


Overall, 372 of 816 participants (45.6%) were hospitalized within 24 hours: 178 of 409 (43.5%) in the magnesium group vs 194 of 407 (47.7%) in the placebo group (difference, -4.2%; absolute risk difference 95% [exact] CI, -11% to 2.8%;  $P = .26$ ; Table 3, Table 4, and Table 5). Specifically, 364 children were hospitalized at the index ED visit (magnesium: 172 [42.1%] vs placebo: 192 [47.2%]) and 8 were admitted during a return visit after initial discharge (magnesium: 6, placebo: 2). Controlling for stratification at randomization for age group and site confirmed lack of statistically significant association between treatment assignment and the primary outcome (Table 3, Table 4, and Table 5). The analysis restricted to the 756 children who completed all experimental treatments also did not reveal a significant association between hospitalization and treatment assignment (Table 3).

There was no statistically significant treatment effect of magnesium and albuterol vs placebo and albuterol in any of the subgroups (Figure 2).

### Secondary Outcomes

There were no significant between-group differences in the changes from baseline to 240 minutes in PRAM score (difference in changes, 0.14 points [95% CI, -0.23 to 0.51];  $P = .45$ ); respiratory rate (0.31 breaths/min [95% CI, -1.17 to 1.79];

Figure 1. Enrollment and Randomization in the Trial



PRAM indicates Pediatric Respiratory Assessment Measure.

<sup>a</sup> Some participants were excluded for more than 1 criterion.

<sup>b</sup> See Table 1 for PRAM calculation and severity interpretation.

<sup>c</sup> Children with previous wheeze not treated with bronchodilators or corticosteroids.

<sup>d</sup> Randomization was stratified by site and age group

Table 2. Characteristics of the Enrolled Participants at Randomization

Characteristic	Group, No. (%)	
	Magnesium (n = 409)	Placebo (n = 407)
Age, median (IQR), y	4.0 (3.0-7.0)	5.0 (3.0-7.0)
Age ≤5 y	253 (61.9)	250 (61.4)
Sex		
Male	257 (62.8)	260 (63.9)
Female	152 (37.2)	147 (36.1)
Hospitalized for asthma in preceding year	105/408 (25.7)	92/401 (22.9)
Previous ICU admission for asthma	43/408 (10.5)	36/401 (9.0)
Personal history of atopy <sup>a</sup>	253/401 (63.1)	240/398 (60.3)
Family history of atopy <sup>b</sup>	307/403 (76.2)	296/398 (74.4)
Preschool wheeze <sup>c</sup>	63/363 (17.4)	56/349 (16.0)
Upper respiratory infection	293 (71.6)	279 (68.6)
Duration of respiratory distress prior to ED arrival, median (IQR), h	16.0 (12.0-24.0)	17.0 (10.0-24.0)
Prior ED visit during this episode	78 (19.1)	79 (19.4)
Self-administered albuterol within 48 h preceding ED arrival	368 (90.0)	374 (91.9)
Oral corticosteroid administered within 48 h preceding ED arrival	73 (17.8)	72 (17.7)
Inhaled corticosteroid administered within 48 h preceding ED arrival	248 (60.6)	223 (54.8)
PRAM score, median (IQR) <sup>d</sup>	6 (5-7)	6 (5-7)
PRAM score ≥8 <sup>d</sup>	61 (14.9)	69 (16.9)
Respiratory rate/min, median (IQR) <sup>d</sup>	36 (32-44)	38 (30-44)
Heart rate/min, median (IQR) <sup>d</sup>	147 (135-158)	146 (134-160)
Oxygen saturation, median (IQR), % <sup>d</sup>	94 (92-96)	94 (92-96)
Blood pressure, median (IQR), mm Hg <sup>d</sup>		
Systolic	108 (101-116)	106 (100-115)
Diastolic	62 (56-68)	63 (57-70)

Abbreviations: ED, emergency department; DBP, diastolic blood pressure; ICU, intensive care unit; IQR, interquartile range; PRAM, Pediatric Respiratory Assessment Measure (see Table 1 for calculation and interpretation); SBP, systolic blood pressure.

<sup>a</sup> History of asthma, allergic rhinitis, or eczema.

<sup>b</sup> History of asthma, allergic rhinitis, or eczema in parents or siblings.

<sup>c</sup> Age 2-5 years, with history of no cough/wheeze between colds and no personal atopy.

<sup>d</sup> Measured at enrollment.

Table 3. Primary Outcome of Hospitalization and Secondary Outcome of Need for Additional Albuterol<sup>a</sup>

Outcome	Group, No. (%)		Unadjusted		Adjusted	
	Magnesium (n = 409)	Placebo (n = 407)	Risk difference (95% CI)	P value	Risk difference (95% CI) <sup>b</sup>	P value
<b>Primary outcome</b>						
Hospitalization within 24 h						
All patients	178 (43.5)	194 (47.7)	-0.04 (-0.11 to 0.03)	.26	-0.05 (-0.13 to 0.02)	.18
Patients with full experimental therapy <sup>c</sup>	158/361 (43.8)	188/395 (47.6)	-0.04 (-0.11 to 0.03)	.29	-0.04 (-0.10 to 0.03)	.25
<b>Secondary outcome (additional albuterol treatments within 240 min)<sup>d</sup></b>						
Additional albuterol treatments, unadjusted mean (95% CI)	1.49 (1.32-1.69)	1.59 (1.41-1.79)	0.94 (0.79 to 1.11) <sup>e</sup>	.47	0.94 (0.78 to 1.14) <sup>e</sup>	.53

<sup>a</sup> All comparisons have controlled for stratification at randomization for site and age group.

<sup>b</sup> Post hoc adjustment was made for stratification at randomization for age group, with site as a random effect.

<sup>c</sup> This was a sensitivity analysis, with the full experimental therapy defined as completion of all 3 study treatments.

<sup>d</sup> The 240-minute interval was chosen to show the potential effect of magnesium over the time period when most hospitalization decisions in asthma are made.

<sup>e</sup> Risk ratio (95% CI).

Table 4. Secondary Outcomes of the Trial

Outcome	Magnesium group (n = 409)			Placebo group (n = 407)			Unadjusted		Adjusted	
	Preintervention, mean (SD)	Postintervention, mean (SD)	Unadjusted difference (95% CI)	Preintervention, mean (SD)	Postintervention, mean (SD)	Unadjusted difference (95% CI)	Difference-in-difference (95% CI)	P value	Difference-in-difference (95% CI) <sup>a</sup>	P value
<b>Changes from baseline to 240 min<sup>b</sup></b>										
PRAM <sup>c</sup>	6.18 (1.33)	3.84 (1.94)	2.43 (2.16 to 2.69)	6.37 (1.27)	4.13 (2.04)	2.29 (2.04 to 2.54)	0.14 (-0.23 to 0.50)	.46	0.14 (-0.23 to 0.51)	.45
Respiratory rate (breaths/min)	38.09 (9.41)	34.35 (8.86)	4.01 (2.93 to 5.09)	38.21 (9.86)	34.54 (8.86)	3.84 (2.81 to 4.87)	0.17 (-1.32 to 1.67)	.82	0.31 (-1.17 to 1.79)	.68
Oxygen saturation (%)	94.00 (3.11)	94.50 (3.04)	-0.91 (-1.27 to -0.55)	94.20 (3.07)	94.90 (2.96)	-0.87 (-1.21 to -0.53)	-0.04 (-0.53 to 0.46)	.88	-0.05 (-0.54 to 0.45)	.86
Systolic blood pressure, mm Hg	108.4 (11.82)	108.31 (12.41)	0.24 (-1.41 to 1.89)	108.0 (10.75)	108.77 (12.96)	-0.54 (-2.08 to 1.01)	0.78 (-1.48 to 3.03)	.50	0.61 (-1.64 to 2.85)	.60
Diastolic blood pressure, mm Hg	62.59 (11.24)	59.64 (11.6)	3.12 (1.41 to 4.83)	63.13 (10.65)	58.85 (10.47)	4.45 (2.84 to 6.07)	-1.33 (-3.68 to 1.02)	.27	-1.35 (-3.70 to 0.99)	.26

Abbreviation: PRAM, Pediatric Respiratory Assessment Measure.

<sup>a</sup> Post hoc adjustment was made for stratification at randomization for age group, with site as a random effect.

<sup>b</sup> The 240-minute interval was chosen to show the potential effect of

magnesium over the time period when most hospitalization decisions in asthma are made. Additional time points are included in eTable 5 in Supplement 5.

<sup>c</sup> See Table 1 for score calculation and severity interpretation.

Table 5. Other Outcomes of the Trial

Outcome	Group, No. (%)		Unadjusted		Adjusted	
	Magnesium (n = 409)	Placebo (n = 407)	Risk difference (95% CI)	P value	Risk difference (95% CI) <sup>a</sup>	P value
Hospitalization within 72 h	180 (44.01)	196 (48.16)	-0.04 (-0.11 to 0.03)	.23	-0.05 (-0.10 to 0.02)	.19
Revisit to any medical facility within 72 h	21/236 (8.90)	15/215 (6.98)	0.02 (-0.03 to 0.07)	.45	0.02 (-0.07 to 0.57)	.54
Intravenous magnesium in emergency department <sup>b</sup>	100 (24.45)	115 (28.26)	-0.04 (-0.10 to 0.02)	.22	-0.04 (-0.09 to 0.02)	.19


<sup>a</sup> Post hoc adjustment was made for stratification at randomization for age group, with site as a random effect.

<sup>b</sup> Intravenous magnesium was administered after the experimental therapy.

$P = .68$ ); oxygen saturation (-0.05% [95% CI, -0.54% to 0.45%];  $P = .86$ ); systolic blood pressure (0.61 mm Hg [95% CI, -1.64 to 2.85];  $P = .60$ ); or mean number of additional albuterol treatments (magnesium: 1.49, placebo: 1.59 [adjusted risk ratio, 0.94 [95% CI, 0.78 to 1.14];  $P = .53$ ). While the small differences in

diastolic blood pressure changes at 20, 40, and 60 minutes reached significance, the directionality was contrary to the study hypothesis. The results of the changes in these outcomes to prespecified end points are summarized in eAppendix 7 in Supplement 5 and eTable 5 in Supplement 5. The

Figure 2. Subgroup Analyses for the Primary Outcome



Numbers and percentages shown are number of patients hospitalized / number of patients at risk. PRAM indicates Pediatric Respiratory Assessment Measure; see Table 1 for details. Preschool wheeze is defined as age 5 years and younger without atopy or cough between upper respiratory infections. Atopy indicates a personal history of asthma, eczema, or allergic rhinitis.

Table 6. Adverse Events in Study Participants

Adverse event <sup>a,b</sup>	Relation to study drug <sup>c</sup>	Magnesium group (n = 409)	Placebo group (n = 407)
Nausea/vomiting	Total	9	5
	Unlikely	5	4
	Possibly	3	1
	Other	1	0
Sore throat/nose, burning tongue, epistaxis	Total	8	0
	Possibly	7	0
	Definitely	1	0
Rash	Total	2	1
	Unlikely	1	1
	Possibly	1	0
Ear pain	Possibly	1	0
Headache	Unlikely	0	1
Hyperglycemia	Unlikely	1	0
Hypertension	Possibly	1	0
Hypotension	Unlikely	0	1
Metabolic acidosis	Unlikely	2	0
Night terrors	Unlikely	0	1
Possible pneumonia	Unlikely	0	1
Possible sepsis	Unlikely	0	1
Status asthmaticus	Unlikely	1	0
Any adverse event		25	11
Any serious adverse event <sup>d</sup>		5	14

<sup>a</sup> No patient had more than 1 adverse event.

<sup>b</sup> These a priori-defined asthma/study-related occurrences were not considered adverse events: cough, respiratory distress (disease-related), asthma-related hospitalization, intravenous insertion, sinus tachycardia, and bitter/salty taste of the experimental solution.

<sup>c</sup> Relation of adverse events to the study drug was evaluated by site investigators according to clinical relationship or theoretical pharmacologic relationship, using the following scale: not related, unlikely related, possibly related, definitely related, other.

<sup>d</sup> Serious adverse events were defined a priori as hypotension requiring medical intervention, apnea, or admission to intensive care unit. All observed serious adverse events consisted of admissions to intensive care unit and none were attributed to the experimental therapy.

number of additional albuterol treatments administered within 240 minutes was also not statistically significantly different between the trial groups (Table 3, Table 4, and Table 5).

**Other Outcomes**

There was no statistically significant treatment effect on hospitalizations or revisits within 72 hours or on administration of IV magnesium after the experimental therapy (Table 3,

Table 4, and Table 5). Adverse events were infrequent and generally not attributed to the experimental therapy (Table 6). A total of 33 of 36 AEs were judged to be mild and 3 were moderate (magnesium: 1 [status asthmaticus]; placebo: 2 [pneumonia, sepsis with antibiotic therapy]). SAEs consisted exclusively of asthma-related admissions to the intensive care unit (Table 6) and none were attributed to the experimental treatment.

## Discussion

In this randomized trial of children with acute asthma refractory to optimized initial therapy, administration of nebulized magnesium with albuterol did not significantly decrease hospitalizations within 24 hours compared with placebo with albuterol.

An important challenge in the management of moderate to severe acute asthma is that the response to the first-line ED treatment is variable. One well-documented reason for these treatment failures are genetic polymorphisms leading to varied response to inhaled  $\beta_2$  agonists and delayed improvement after corticosteroids.<sup>33-36</sup> In children refractory to the initial optimized therapy, inhaled magnesium would in theory seem to be an attractive second-line agent. To date, however, there have been relatively few published studies, and those have been criticized for use of poorly standardized entry criteria, heterogeneous outcome measures, and widely varying co-interventions.<sup>8,37</sup> Consequently, current systematic reviews of nebulized magnesium in childhood asthma conclude there is substantial uncertainty about its effectiveness.<sup>8,37,38</sup>

More specifically, 2 larger pediatric randomized trials of inhaled magnesium have been published.<sup>14,15</sup> On one hand, Alansari et al<sup>15</sup> concluded that nebulized magnesium did not significantly reduce time to discharge readiness in 365 hospitalized patients in a single center in Qatar. On the other hand, in a multicenter ED-based trial of 508 children in the United Kingdom, Powell et al<sup>14</sup> demonstrated that receipt of inhaled magnesium resulted in a statistically significant but clinically small improvement in the Yung asthma score. This study led the British Thoracic Society Scottish Intercollegiate Guidelines Network to recommend that inhaled magnesium be added to each bronchodilator treatment in the first hour in children with severe asthma and short symptom duration or oxygen saturation of less than 92%.<sup>39</sup>

To overcome limitations of previous studies, this trial used standardized patient entry criteria assessed by a validated PRAM score, optimized initial therapy prior to screening for study eligibility, tested a large magnesium dose delivered via high-efficiency nebulizer, and ensured adequate power for assessment of a health care utilization outcome. This study provides evidence that children who remain in moderate to severe respiratory distress despite aggressive initial therapy do not derive benefit from inhaled magnesium.

Plausible reasons for lack of benefit of inhaled magnesium deserve consideration. First, it is unlikely that insufficient magnesium dose or inadequate delivery were responsible. While 7% of children did not receive all 3 magnesium doses, analyses limited to children who received all 3 doses showed similar results, suggesting that this issue did not confound the results. Second, because a high-efficiency nebulizer was used to administer albuterol along with the study drug

and/or placebo, participants received a much larger dose of albuterol to the lower airways than is achieved with conventional nebulizers. Therefore, some children who experienced inadequate responses to the initial treatment regimen, which included standard dose of albuterol, may have improved due to the augmented albuterol deposition provided by the high-efficiency nebulizer. This response may have masked any added benefit of magnesium. Additionally, while the pragmatic study design may have increased generalizability of the study findings, lack of standardized hospitalization criteria may have contributed to the null result.

## Limitations

This trial has several limitations. First, it was not feasible to confirm physician-diagnosed asthma at the time of enrollment in all participants. To mitigate the risk of enrollment of children without asthma, children younger than 2 years of age and those presenting with their first episode of asthmalike phenotype who may have had other diagnoses were excluded.

Second, only a small proportion of enrolled children had severe respiratory distress because some children with very high presenting PRAM scores received IV magnesium shortly after arrival based on physician judgment.

Third, while severe PRAM denotes a risk for hospitalization, the subgroup analyses suggest lack of statistically significant treatment effect of nebulized magnesium in children with PRAM scores of 8 or higher. Therefore, the results do not support that children with severe presentation might derive benefit from magnesium.

Fourth, while the PRAM scoring has been shown to have good interrater reliability<sup>20</sup> and PRAM is routinely used to monitor response to therapy in children with asthma in Canadian EDs, the decision to hospitalize children is not standardized and is influenced by multiple patient-level and system factors. However, the randomized study design used does minimize the effect of such variation on the outcomes of interest.

Fifth, because the 95% CI around the difference in the primary outcome includes 10%, the study may have been underpowered to detect small differences in favor of magnesium. In addition, the study results are not generalizable to magnesium therapy administered intravenously.

## Conclusions

Among children with refractory acute asthma in the ED, nebulized magnesium with albuterol, compared with placebo with albuterol, did not significantly decrease the hospitalization rate for asthma within 24 hours. The findings do not support use of nebulized magnesium with albuterol among children with refractory acute asthma.

### ARTICLE INFORMATION

**Accepted for Publication:** September 21, 2020.

**Author Affiliations:** Division of Pediatric Emergency Medicine, The Hospital for Sick

Children, University of Toronto, Toronto, Ontario, Canada (Schuh, Sweeney, Rumantir); Research Institute, The Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada (Schuh, Finkelstein); Division of Respiratory Medicine,

Research Institute, The Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada (Coates); Ontario Child Health Support Unit, SickKids Research Institute, Toronto, Ontario, Canada (Willan); Dalla Lana School of Public Health,



University of Toronto, Toronto, Ontario, Canada (Willan); Research Institute, Clinical Research Services, The Hospital for Sick Children, Toronto, Ontario, Canada (Stephens); Department of Biostatistics, Princess Margaret Cancer Centre, University of Toronto, Toronto, Ontario, Canada (Atenafu); Divisions of Pediatric Emergency Medicine and Clinical Pharmacology and Toxicology, The Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada (Finkelstein); Departments of Pediatrics and Emergency Medicine, Alberta Children's Hospital, University of Calgary, Calgary, Alberta, Canada (Thompson); Alberta Children's Hospital Research Institute, University of Calgary, Calgary, Alberta, Canada (Thompson); Department of Pediatrics and Emergency Medicine, Children's Hospital of Eastern Ontario, University of Ottawa, Ottawa, Ontario, Canada (Zemek, Plint); CHEO Research Institute, Ottawa, Ontario, Canada (Zemek, Plint); Department of Pediatric Emergency Medicine, CHU Sainte-Justine, Montréal, Quebec, Canada (Gravel, Ducharme); Department of Pediatrics and of Social and Preventive Medicine, Université de Montréal, Montréal, Quebec, Canada (Gravel, Ducharme); Departments of Pediatrics, Emergency Medicine, and Physiology & Pharmacology, Alberta Children's Hospital, Calgary, Alberta, Canada (Johnson); Alberta Children's Hospital Research Institute, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada (Johnson); Division of Pediatric Emergency Medicine, British Columbia Children's Hospital, University of British Columbia, Vancouver, British Columbia, Canada (Black); Departments of Pediatrics and Emergency Medicine, Stollery Children's Hospital, University of Alberta, Edmonton, Alberta, Canada (Curtis); Pediatrics/Pediatric Emergency Medicine, University of Manitoba, Winnipeg, Manitoba, Canada (Beer); Children's Hospital Research Institute of Manitoba, Winnipeg, Manitoba, Canada (Beer); Department of Pediatrics and Child Health, University of Manitoba, Winnipeg, Manitoba, Canada (Klassen); Children's Hospital Research Institute of Manitoba, Winnipeg, Manitoba, Canada (Klassen); Pharmacy Department, The Hospital for Sick Children, Toronto, Ontario, Canada (Nicksy); Sections of Pediatric Emergency Medicine and Gastroenterology, Departments of Pediatrics and Emergency Medicine, Alberta Children's Hospital, Alberta Children's Hospital Research Institute, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada (Freedman).

**Author Contributions:** Dr Schuh had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** Schuh, Coates, Willan, Zemek, Plint, Gravel, Ducharme, Johnson, Black, Curtis, Klassen, Nicksy, Freedman.

**Acquisition, analysis, or interpretation of data:** Sweeney, Rumantir, Willan, Stephens, Atenafu, Finkelstein, Thompson, Zemek, Plint, Gravel, Black, Curtis, Beer, Nicksy, Freedman.

**Drafting of the manuscript:** Schuh, Stephens, Curtis, Nicksy, Freedman.

**Critical revision of the manuscript for important intellectual content:** Schuh, Sweeney, Rumantir, Coates, Willan, Stephens, Atenafu, Finkelstein, Thompson, Zemek, Plint, Gravel, Ducharme, Johnson, Black, Curtis, Beer, Klassen, Freedman.

**Statistical analysis:** Willan, Stephens, Atenafu.

**Obtained funding:** Schuh, Zemek, Plint, Gravel, Johnson, Curtis, Freedman.

**Administrative, technical, or material support:** Sweeney, Rumantir, Coates, Finkelstein, Thompson, Gravel, Ducharme, Black, Beer, Nicksy, Freedman. **Supervision:** Schuh, Sweeney, Rumantir, Johnson, Beer, Klassen.

**Conflict of Interest Disclosures:** Dr Schuh reported receiving grants from the Canadian Institutes for Health Research (CIHR), Thrasher Research Fund, Physicians' Services Incorporated Foundation, and Hospital for Sick Children and nonfinancial support from La Diffusion Technique Francaise during the conduct of the study. Dr Sweeney reported receiving grants from the Hospital for Sick Children during the conduct of the study. Dr Finkelstein reported receiving grants from CIHR during the conduct of the study. Dr Zemek reported holding competitively funded research grants from CIHR, Ontario Neurotrauma Foundation, Physicians' Services Incorporated Foundation, CHEO Foundation, Ontario Brain Institute, Ontario SPOR Support Unit, National Football League, and Clinical Research Chair in Pediatric Concussion from the University of Ottawa. He has no commercial conflicts of interest to disclose. Dr Gravel reported receiving grants from CIHR during the conduct of the study. Dr Ducharme reported receiving unrestricted research funds from GlaxoSmithKline, AstraZeneca, Novartis, Teva, and Trudell Medical; grants from GlaxoSmithKline, Covis, Thorasys, and CIHR; and personal fees from Thorasys, Covis, and Teva outside the submitted work. No other disclosures were reported.

**Funding/Support:** This research was supported by grants from CIHR, the Thrasher Research Fund, the Physicians' Services Incorporated Foundation, and the Hospital for Sick Children. Dr Freedman is supported by the Alberta Children's Hospital Foundation Professorship in Child Health and Wellness.

**Role of the Funder/Sponsor:** The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Group Information:** The Pediatric Emergency Research Canada (PERC) Network members include the following: Suzanne Schuh, MD, Judy Sweeney, RN, BScN, Maggie Rumantir, MD, Allan L. Coates, MDCM, BEng, Andrew R. Willan, PhD, Derek Stephens, MSc, BSc, Yaron Finkelstein, MD, and Darcy Nicksy, BScPhM (The Hospital for Sick Children, Toronto, Ontario, Canada); Graham Thompson, MD, and Stephen B. Freedman, MDCM, MSc (Alberta Children's Hospital, Calgary, Alberta, Canada); Darcy Beer, MD, and Terry Klassen, MD, MSc (Winnipeg Children's Hospital, Winnipeg, Manitoba, Canada); Sarah Curtis, MD (Stollery Children's Hospital, Edmonton, Alberta, Canada); Jocelyn Gravel, MD, MSc, and Francine Ducharme, MD, MSc (Sainte-Justine Pédiatrie, Montreal, Quebec, Canada); Roger Zemek, MD, and Amy Plint, MD, MSc (Children's Hospital of Eastern Ontario, Ottawa, Ontario, Canada); and Karen Black, MD, MSc (British Columbia Children's Hospital, Vancouver, British Columbia, Canada).

**Meeting Presentation:** This study was accepted for a platform presentation at the annual conference of the Pediatric Academic Societies; May 2020; Philadelphia, Pennsylvania.

**Additional Contributions:** We thank the participants and their families for trusting us to conduct this trial; the trial investigators and support staff across all sites for their commitment to the successful conduct of the trial; the site coordinators, including Rena Papadimitropoulos (research coordinator, Children's Hospital of Winnipeg), Arpita Majumdar (MSc, Children's Hospital of Winnipeg), Jianling Xie (MD, MPH, Alberta Children's Hospital), Dale Dagleish (RN, BHSn, CHEO), Candice McGahern (BA, CHEO), Ally Slattery (RN, British Columbia Children's Hospital), Maryse Lagace (RN, BScN, CCRP, CHU Sainte-Justine, Montreal), Manasi Rajagopal (research coordinator, Stollery Children's Hospital), and Nadia Dow (research coordinator, Stollery Children's Hospital) for trial coordination; the emergency department physicians, nurses, and ancillary staff at all sites; the staff of the Pediatric Research Academic Initiative at SickKids Emergency (PRAISE) and of the SickKids Emergency Assistants for Research in Child Health (SEARCH) programs for identifying potentially eligible participants; Tanveer H. Collins (MD, The Hospital for Sick Children) for his help with the database; Aya Finkelstein (The Hospital for Sick Children) for her help with the study database; and Lejla Halilovic (BSc, The Hospital for Sick Children) for her administrative assistance with the manuscript preparation. The site coordinators and Dr Collins received financial compensation for their role in the study. The other contributors were not compensated.

**Data Sharing Statement:** See Supplement 6.

## REFERENCES

- Moorman JE, Rudd RA, Johnson CA, et al; Centers for Disease Control and Prevention (CDC). National surveillance for asthma: United States, 1980-2004. *MMWR Surveill Summ*. 2007;56(8):1-54.
- National Asthma Education and Prevention Program. Expert Panel Report 3 (EPR-3): guidelines for the diagnosis and management of asthma: summary report 2007. *J Allergy Clin Immunol*. 2007;120(5)(suppl):S94-S138. doi:10.1016/j.jaci.2007.09.029
- Global Initiative for Asthma. Global Initiative for Asthma (GINA). Accessed April 5, 2020. <http://www.ginasthma.org>
- Ortiz-Alvarez O, Mikrogianakis A; Canadian Paediatric Society, Acute Care Committee. Managing the paediatric patient with an acute asthma exacerbation. *Paediatr Child Health*. 2012;17(5):251-262. doi:10.1093/pch/17.5.251
- Tse SM, Tantisira K, Weiss ST. The pharmacogenetics and pharmacogenomics of asthma therapy. *Pharmacogenomics J*. 2011;11(6):383-392. doi:10.1038/tj.2011.46
- Cheuk DK, Chau TC, Lee SL. A meta-analysis on intravenous magnesium sulphate for treating acute asthma. *Arch Dis Child*. 2005;90(1):74-77. doi:10.1136/adc.2004.050005
- Griffiths B, Kew KM. Intravenous magnesium sulfate for treating children with acute asthma in the emergency department. *Cochrane Database Syst Rev*. 2016;4:CD011050.
- Su Z, Li R, Gai Z. Intravenous and nebulized magnesium sulfate for treating acute asthma in children: a systematic review and meta-analysis.

- Pediatr Emerg Care.* 2018;34(6):390-395. doi:10.1097/PEC.0000000000000909
9. Liu X, Yu T, Rower JE, Campbell SC, Sherwin CM, Johnson MD. Optimizing the use of intravenous magnesium sulfate for acute asthma treatment in children. *Pediatr Pulmonol.* 2016;51(12):1414-1421. doi:10.1002/ppul.23482
  10. Schuh S, Macias C, Freedman SB, et al. North American practice patterns of intravenous magnesium therapy in severe acute asthma in children. *Acad Emerg Med.* 2010;17(11):1189-1196. doi:10.1111/j.1553-2712.2010.00913.x
  11. Johnson MD, Zorc JJ, Nelson DS, et al; Pediatric Emergency Care Applied Research Network (PECARN). Intravenous magnesium in asthma pharmacotherapy: variability in use in the PECARN Registry. *J Pediatr.* 2020;220:165-174.e2. doi:10.1016/j.jpeds.2020.01.062
  12. Mahajan P, Haritos D, Rosenberg N, Thomas R. Comparison of nebulized magnesium sulfate plus albuterol to nebulized albuterol plus saline in children with acute exacerbations of mild to moderate asthma. *J Emerg Med.* 2004;27(1):21-25. doi:10.1016/j.jemermed.2004.02.006
  13. Meral A, Coker M, Tanaç R. Inhalation therapy with magnesium sulfate and salbutamol sulfate in bronchial asthma. *Turk J Pediatr.* 1996;38(2):169-175.
  14. Powell C, Kolamunnage-Dona R, Lowe J, et al; MAGNETIC study group. Magnesium sulphate in acute severe asthma in children (MAGNETIC): a randomised, placebo-controlled trial. *Lancet Respir Med.* 2013;1(4):301-308. doi:10.1016/S2213-2600(13)70037-7
  15. Alansari K, Ahmed W, Davidson BL, Alamri M, Zakaria I, Alrifai M. Nebulized magnesium for moderate and severe pediatric asthma: a randomized trial. *Pediatr Pulmonol.* 2015;50(12):1191-1199. doi:10.1002/ppul.23158
  16. Schuh S, Sweeney J, Freedman SB, et al; Pediatric Emergency Research Canada Group. Magnesium nebulization utilization in management of pediatric asthma (MagNUM PA) trial: study protocol for a randomized controlled trial. *Trials.* 2016;17(1):261. doi:10.1186/s13063-015-1151-x
  17. Ducharme FM, Chalut D, Plotnick L, et al. The Pediatric Respiratory Assessment Measure: a valid clinical score for assessing acute asthma severity from toddlers to teenagers. *J Pediatr.* 2008;152(4):476-480, 480.e1. doi:10.1016/j.jpeds.2007.08.034
  18. US Committee on Extension to the Standard Atmosphere. *US Standard Atmosphere, 1976.* US Government Printing Office; 1976.
  19. Bhogal SK, McGillivray D, Bourbeau J, Benedetti A, Bartlett S, Ducharme FM. Early administration of systemic corticosteroids reduces hospital admission rates for children with moderate and severe asthma exacerbation. *Ann Emerg Med.* 2012; 60(1):84-91.e3. doi:10.1016/j.annemergmed.2011.12.027
  20. Chalut DS, Ducharme FM, Davis GM. The Preschool Respiratory Assessment Measure (PRAM): a responsive index of acute asthma severity. *J Pediatr.* 2000;137(6):762-768. doi:10.1067/mpd.2000.110121
  21. Birken CS, Parkin PC, Macarthur C. Asthma severity scores for preschoolers displayed weaknesses in reliability, validity, and responsiveness. *J Clin Epidemiol.* 2004;57(11):1177-1181. doi:10.1016/j.jclinepi.2004.02.016
  22. Lehr AR, McKinney ML, Gouin S, Blais JG, Pusic MV, Ducharme FM. Development and pretesting of an electronic learning module to train health care professionals on the use of the Pediatric Respiratory Assessment Measure to assess acute asthma severity. *Can Respir J.* 2013;20(6):435-441. doi:10.1155/2013/148645
  23. Aggarwal P, Sharad S, Handa R, Dwiwedi SN, Irshad M. Comparison of nebulized magnesium sulphate and salbutamol combined with salbutamol alone in the treatment of acute bronchial asthma: a randomised study. *Emerg Med J.* 2006;23(5):358-362. doi:10.1136/emj.2005.026203
  24. Tal A, Golan H, Grauer N, Aviram M, Albin D, Quastel MR. Deposition pattern of radiolabeled salbutamol inhaled from a metered-dose inhaler by means of a spacer with mask in young children with airway obstruction. *J Pediatr.* 1996;128(4):479-484. doi:10.1016/S0022-3476(96)70357-8
  25. Coates AL, Leung K, Vecellio L, Schuh S. Testing of nebulizers for delivering magnesium sulfate to pediatric asthma patients in the emergency department. *Respir Care.* 2011;56(3):314-318. doi:10.4187/respcare.00826
  26. Coates AL, Leung K, Chan J, Ribeiro N, Charron M, Schuh S. Respiratory system deposition with a novel aerosol delivery system in spontaneously breathing healthy adults. *Respir Care.* 2013;58(12):2087-2092. doi:10.4187/respcare.02455
  27. Ungar WJ, Coyte PC; Pharmacy Medication Monitoring Program Advisory Board. Prospective study of the patient-level cost of asthma care in children. *Pediatr Pulmonol.* 2001;32(2):101-108. doi:10.1002/ppul.1095
  28. Plotnick LH, Ducharme FM. Combined inhaled anticholinergic agents and beta-2-agonists for initial treatment of acute asthma in children. [Update in *Cochrane Database Syst Rev.* 2000;(4): CD000060]. *Cochrane Database Syst Rev.* 2000; (2):CD000060. doi:10.1002/14651858.CD000060
  29. Holm S. A simple sequentially rejective multiple test procedure. *Scand J Stat.* 1979:65-70.
  30. Centers for Disease Control and Prevention (CDC). Asthma mortality and hospitalization among children and young adults: United States, 1980-1993. *MMWR Morb Mortal Wkly Rep.* 1996;45(17):350-353.
  31. Bjerg A, Sandström T, Lundbäck B, Rönmark E. Time trends in asthma and wheeze in Swedish children 1996-2006: prevalence and risk factors by sex. *Allergy.* 2010;65(1):48-55. doi:10.1111/j.1398-9995.2009.02105.x
  32. Castro-Rodríguez JA, Holberg CJ, Wright AL, Martinez FD. A clinical index to define risk of asthma in young children with recurrent wheezing. *Am J Respir Crit Care Med.* 2000;162(4, pt 1):1403-1406. doi:10.1164/ajrccm.162.4.991211
  33. Carroll CL, Sala KA, Zucker AR, Schramm CM. Beta-adrenergic receptor polymorphisms associated with length of ICU stay in pediatric status asthmaticus. *Pediatr Pulmonol.* 2012;47(3):233-239. doi:10.1002/ppul.21544
  34. Chun E, Lee HS, Bang BR, et al. Dexamethasone-induced FKBP51 expression in peripheral blood mononuclear cells could play a role in predicting the response of asthmatics to treatment with corticosteroids. *J Clin Immunol.* 2011;31(1):122-127. doi:10.1007/s10875-010-9463-9
  35. Lima JJ. Do genetic polymorphisms alter patient response to inhaled bronchodilators? *Expert Opin Drug Metab Toxicol.* 2014;10(9):1231-1240. doi:10.1517/17425255.2014.939956
  36. Moore PE, Ryckman KK, Williams SM, Patel N, Summar ML, Sheller JR. Genetic variants of GSNOR and ADRB2 influence response to albuterol in African-American children with severe asthma. *Pediatr Pulmonol.* 2009;44(7):649-654. doi:10.1002/ppul.21033
  37. Normansell R, Knightly R, Milan SJ, Knopp-Sihota JA, Rowe BH, Powell C. Inhaled magnesium sulfate in the treatment of acute asthma in children. *Paediatr Respir Rev.* 2018;26:31-33. doi:10.1016/j.prrv.2018.01.001
  38. Knightly R, Milan SJ, Hughes R, et al. Inhaled magnesium sulfate in the treatment of acute asthma. *Cochrane Database Syst Rev.* 2017;11: CD003898. doi:10.1002/14651858.CD003898.pub6
  39. British Thoracic Society; Scottish Intercollegiate Guidelines Network. British guideline on the management of asthma. *Thorax.* 2014;69(suppl 1):i-192.