

# Pediatric In-Hospital Cardiac Arrest and Cardiopulmonary Resuscitation in the United States

## A Review

Ryan W. Morgan, MD, MTR; Matthew P. Kirschen, MD, PhD; Todd J. Kilbaugh, MD; Robert M. Sutton, MD, MSCE; Alexis A. Topjian, MD, MSCE

**IMPORTANCE** Pediatric in-hospital cardiac arrest (IHCA) occurs frequently and is associated with high morbidity and mortality. The objective of this narrative review is to summarize the current knowledge and recommendations regarding pediatric IHCA and cardiopulmonary resuscitation (CPR).

**OBSERVATIONS** Each year, more than 15 000 children receive CPR for cardiac arrest during hospitalization in the United States. As many as 80% to 90% survive the event, but most patients do not survive to hospital discharge. Most IHCAs occur in intensive care units and other monitored settings and are associated with respiratory failure or shock. Bradycardia with poor perfusion is the initial rhythm in half of CPR events, and only about 10% of events have an initial shockable rhythm. Pre-cardiac arrest systems focus on identifying at-risk patients and ensuring that they are in monitored settings. Important components of CPR include high-quality chest compressions, timely defibrillation when indicated, appropriate ventilation and airway management, administration of epinephrine to increase coronary perfusion pressure, and treatment of the underlying cause of cardiac arrest. Extracorporeal CPR and measurement of physiological parameters are evolving areas in improving outcomes. Structured post-cardiac arrest care focused on targeted temperature management, optimization of hemodynamics, and careful intensive care unit management is associated with improved survival and neurological outcomes.

**CONCLUSIONS AND RELEVANCE** Pediatric IHCA occurs frequently and has a high mortality rate. Early identification of risk, prevention, delivery of high-quality CPR, and post-cardiac arrest care can maximize the chances of achieving favorable outcomes. More research in this field is warranted.

*JAMA Pediatr.* doi:10.1001/jamapediatrics.2020.5039  
Published online November 23, 2020.

**Author Affiliations:** Department of Anesthesiology and Critical Care Medicine, Division of Critical Care Medicine, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania (Morgan, Kirschen, Kilbaugh, Sutton, Topjian); Department of Anesthesiology and Critical Care Medicine, Perelman School of Medicine at the University of Pennsylvania, Philadelphia (Morgan, Kirschen, Kilbaugh, Sutton, Topjian).

**Corresponding Author:** Ryan W. Morgan, MD, MTR, Division of Critical Care Medicine, Children's Hospital of Philadelphia, 3401 Civic Center Blvd, 6 Wood-6104, Philadelphia, PA 19104 (morganr1@email.chop.edu).

An estimated 15 200 children receive cardiopulmonary resuscitation (CPR) for in-hospital cardiac arrest (IHCA) annually in the United States.<sup>1</sup> Pediatric IHCA outcomes improved during the first decade of the 21st century,<sup>2,3</sup> likely owing to a widespread recognition of the need for high-quality resuscitation and a focus on recognizing and transferring the deteriorating pediatric inpatient to an intensive care unit (ICU). However, survival rates have plateaued and more than half of children with IHCA do not survive to hospital discharge.<sup>3</sup>

Compared with adults with IHCA, children with IHCA differ in demographic characteristics, location within the hospital, cause of cardiac arrest, initial cardiac arrest rhythm, and type of monitoring in place.<sup>4-7</sup> Table 1 shows a comparison of these variables between pediatric and adult patients.<sup>3-8</sup> However, prospective pediatric IHCA studies are limited, and most pediatric resuscitation guidelines are extrapolated from adult guidelines.<sup>9,10</sup> Substantial differences also exist between pediatric IHCA and pediatric out-of-hospital cardiac arrest (OHCA). Although IHCA occurrence is frequently associated with the progression of underlying disease processes, such as shock

and respiratory failure, common causes of pediatric OHCA include drowning, sudden infant death syndrome, and sudden cardiac death from arrhythmias.<sup>11</sup> Compared with IHCA, OHCA survival outcomes are worse,<sup>12</sup> and OHCA systems of care are focused on encouraging bystander CPR and optimizing basic life support measures in the prehospital state.<sup>13</sup>

Pediatric IHCA is a modifiable disease process. Efforts to prevent its occurrence, treat it with high-quality CPR, and deliver successful post-cardiac arrest care can demonstrably improve outcomes. In this narrative review, we discuss pediatric IHCA, including incidence, causes, prevention, treatment, and post-cardiac arrest therapies.

## Methods

This narrative review was based on several informal searches of PubMed for peer-reviewed English-language articles related to pediatric IHCA. The searches were conducted between January 4,

**Table 1. Characteristics of Pediatric vs Adult In-Hospital Cardiac Arrest**

Characteristic	Pediatric patients <sup>3-5,7</sup>	Adult patients <sup>4,6,7</sup>
Incidence	Approximately 15 200/y in United States	Approximately 292 000/y in United States
Age	Median: 1-2 y; mean: 3-5 y <sup>a</sup>	Mean: 66 y
Male sex	50%-55%	60%
Hospital location	85%-90% In monitored settings <sup>b</sup>	50% In monitored settings <sup>b</sup>
Airway in place	Approximately 80%	Approximately 30%-35%
Cause of cardiac arrest	Most with respiratory failure and/or progressive shock	50%-60% Cardiac; 15%-40% respiratory
Initial cardiac arrest rhythm	50% Bradycardia with poor perfusion; 40% other nonshockable rhythms	80% Nonshockable rhythms (pulseless electrical activity or asystole)
Survival to discharge	45%-50% (40% Among those with pulseless events)	25%
Neurological outcome	Favorable in 80%-90% of survivors <sup>c</sup>	Good functional status in 80% of survivors <sup>d</sup>

<sup>a</sup> Age depends on inclusion criteria of studies.

<sup>b</sup> Monitored settings include intensive care units, emergency departments, and operating rooms.

<sup>c</sup> Pediatric Cerebral Performance Category scale (range: 1-6, with 1 indicating normal function and 6 indicating brain death<sup>8</sup>): 1 to 3 or unchanged from baseline.

<sup>d</sup> Cerebral Performance Category scale: 1 or 2.

2020, and April 15, 2020. Searches were limited to studies that included children younger than 18 years. Search terms were as follows: in-hospital cardiac arrest, cardiac arrest, cardiopulmonary resuscitation, resuscitation, rapid response, medical emergency team, post-arrest, and post-resuscitation. We placed no limit on the date ranges of reviewed articles. Because of the small number of prospective randomized trials in pediatric resuscitation, we did not perform a systematic review. Our objective was to review the current knowledge and recommendations regarding pediatric IHCA and CPR.

## Discussion

### Epidemiological Characteristics of Pediatric IHCA

#### Incidence and Location

Because of the lack of a central reporting process, the actual incidence of pediatric IHCA in the United States remains unknown. Currently accepted estimates are based on data from the American Heart Association (AHA) Get With The Guidelines—Resuscitation (GWTG-R) registry. Given the voluntary nature of this prospective quality improvement registry, data from participating hospitals may not be entirely representative of other hospitals. Regardless, every year, an estimated 15 200 children have IHCAs in the United States.<sup>1</sup> This number includes 7100 pulseless IHCAs and 8100 cases of CPR provided for bradycardia with poor perfusion.<sup>1</sup> Including recurrent events, the total annual number of pediatric IHCA cases is approximately 19 900.<sup>1</sup> In accordance with Utstein CPR reporting guidelines,<sup>14</sup> we refer to all CPR events (pulseless rhythms or bradycardia with poor perfusion) as IHCA.

As opposed to adult IHCAs, which are generally divided between monitored and unmonitored in-hospital settings,<sup>7</sup> 85% to 90% of pediatric IHCAs have occurred in monitored settings, most commonly the ICU.<sup>15</sup> This proportion has increased over time,<sup>15</sup> likely because of widespread implementation of rapid response systems

and the recognition and timely transfer of deteriorating patients.<sup>16-19</sup> The incidence of IHCA among children admitted to pediatric ICUs is approximately 1.8%.<sup>5</sup> Children with heart disease have a substantially higher risk of IHCA,<sup>20</sup> and 3.1% of pediatric patients in cardiac ICUs receive CPR during their admission.<sup>21</sup>

#### Causes and Rhythms

Pediatric IHCA most often occurs in the setting of progressive respiratory failure or shock.<sup>4,5,22</sup> Interventions in place at the onset of CPR include invasive mechanical ventilation in 59% to 82% of children and vasopressor infusions in 29% to 78% of children.<sup>23,24</sup>

Most initial cardiac arrest rhythms are nonshockable (ie, asystole, pulseless electrical activity [PEA], or bradycardia with poor perfusion). Bradycardia with poor perfusion is the initial CPR rhythm in more than half of IHCA events. Children with bradycardia with poor perfusion have superior survival outcomes vs survival outcomes in those with pulselessness at CPR onset.<sup>24,25</sup> However, between 31% and 51% of patients with bradycardia and poor perfusion become pulseless during CPR, and these patients have worse outcomes than either those who were initially pulseless or those who were never pulseless.<sup>24,26</sup>

Ventricular fibrillation (VF) and pulseless ventricular tachycardia (PVT) account for approximately 10% of CPR events.<sup>2,4,27</sup> Primary VF and PVT are associated with more favorable outcomes than PEA and asystole. However, when they occur as secondary rhythms during CPR (after initial PEA or asystole), the outcomes are worse than in those with either primary VF or PVT or in those without VF or PVT at any time.<sup>4,27</sup> This observation may reflect that secondary VF and PVT occur because of worsening myocardial injury over the course of CPR or that secondary VF and PVT are not immediately recognized and thus defibrillated.<sup>27</sup>

#### Outcomes

Survival outcomes are associated with the patient's underlying diagnoses and the cause of the event,<sup>4</sup> the initial cardiac arrest rhythm,<sup>4,27</sup> the quality of CPR delivered and hemodynamics achieved,<sup>23,28,29</sup> the duration of CPR,<sup>5,22</sup> and the quality of post-cardiac arrest care.<sup>30</sup> Although the incidence of IHCA in children with heart disease is relatively high, the survival rates among these patients, particularly those undergoing surgical cardiac treatment during their hospitalization, are higher than among patients without heart disease.<sup>22</sup>

In ICUs, rates of acute IHCA event survival (return of spontaneous circulation [ROSC] or return of circulation with extracorporeal support) are 78% to 90% in recent multicenter observational studies.<sup>5,23</sup> These rates have doubled over the past 2 decades; however, many children with event survival die before hospital discharge.<sup>2,3</sup> According to a recent GWTG-R study, 32% of children with pulseless cardiac arrests from 2000 to 2018 survived to hospital discharge, increasing from 19% in 2000 to 40% in 2011 and then remaining relatively stable through the subsequent 7 years.<sup>3</sup> Among patients with bradycardia with poor perfusion who never became pulseless during their CPR event, 66% survived to hospital discharge in 2018.<sup>3</sup>

Of those patients who survive to discharge, approximately 90% have grossly favorable neurological outcomes,<sup>5,23</sup> as ascertained with the Pediatric Cerebral Performance Category scale (range: 1-6, with 1 indicating normal function and 6 indicating brain death).<sup>8</sup> How-

ever, this scale, typically extrapolated retrospectively from medical record review, lacks sensitivity for more granular neurological dysfunction, and many IHCA event survivors have subtle and sustained neurological impairment.<sup>5,31</sup> Moreover, postresuscitation death is commonly attributed to declaration of death by neurological criteria or poor neurological outcome, which prompts the withdrawal of life-sustaining therapies.<sup>32</sup> Further studies are necessary to better characterize neurocognitive and neurobehavioral outcomes among survivors.

### Prevention of Pediatric IHCA

Rapid response teams and medical emergency teams have become a nearly universal presence in pediatric hospitals over the past 2 decades. Although their composition and characteristics vary,<sup>17</sup> their implementation has been temporally associated with decreases in cardiac arrest frequency and improvements in mortality.<sup>18,33,34</sup> In addition, severity-of-illness scoring systems have been used to identify patients at risk for clinical deterioration but have not consistently been associated with decreased hospital mortality.<sup>35</sup>

Among children in the ICU, previous work has sought to identify those at risk for IHCA to provide situational awareness and facilitate focused treatment to prevent deterioration.<sup>36,37</sup> Across 15 cardiac ICUs, implementation of an IHCA prevention bundle was associated with a decreased incidence of IHCA.<sup>38</sup> In addition, evidence showed that specific ICU staffing models, such as 24 hours per day for 7 days per week in-hospital attending-intensivist coverage, were associated with IHCA prevention.<sup>39</sup>

### Treatment of Cardiac Arrest

The cornerstone of cardiac arrest therapy is the delivery of high-quality CPR that temporarily supports organ perfusion while optimizing the probability of attaining ROSC. The components of CPR and other therapies delivered during IHCA are discussed in this section.

#### Chest Compressions

High-quality CPR consists of chest compressions delivered at an appropriate rate with adequate depth, full chest wall recoil, and minimal interruptions. Compliance with the AHA pediatric basic and advanced life support guidelines recommendations for chest compressions<sup>9,10</sup> is associated with higher rates of survival.<sup>28,29</sup> Table 2 shows the AHA guidelines target for each metric of high-quality CPR.<sup>9,10</sup> Widespread technology is now available, largely in the form of CPR quality-monitoring defibrillators, that measures chest compression mechanics and provides feedback to clinicians in real time. Systematic use of such technology coupled with resuscitation debriefing programs focused on CPR quality is associated with improved outcomes.<sup>40</sup>

Training and equipping clinicians to deliver high-quality CPR according to these metrics has likely been associated with improvement in IHCA outcomes.<sup>2</sup> However, specific targets for pediatric chest compression rates and depths are largely based on expert consensus and extrapolation from adult studies.<sup>10</sup> Rates outside of those recommended by the AHA guidelines have recently been associated with improved outcomes.<sup>41</sup> Moreover, depths within the target range are frequently difficult to achieve, even in high-performing centers that largely achieve other CPR metrics.<sup>42</sup> These

**Table 2. Guideline Targets for Pediatric Cardiopulmonary Resuscitation Quality**

Metric	Target <sup>9,10</sup>
Rate	100-120 Compressions/min
Depth	Adolescents <sup>a</sup> : 5-6 cm
	Children <sup>b</sup> : approximately 5 cm (≥one-third anterior-posterior chest depth)
	Infants <sup>c</sup> : approximately 4 cm (≥one-third anterior-posterior chest depth)
Chest compression fraction	Chest compressions for ≥80% of time of arrest
Chest recoil	Allow full recoil between compressions; avoid leaning
Ventilations	Advanced airway: 10 breaths/min
	No advanced airway (adolescents <sup>a</sup> ): 30:2 compression to ventilation ratio
	No advanced airway (children <sup>b</sup> and infants <sup>c</sup> ): 15:2 compression to ventilation ratio

<sup>a</sup> Adolescents are those showing signs of puberty (eg, breast development in girls; axillary hair in boys).

<sup>b</sup> Children are those 1 year or older without signs of puberty.

<sup>c</sup> Infants are those younger than 1 year.

data highlight the need for further studies to refine the specific targets for chest compression mechanics.

#### Defibrillation

Prompt defibrillation is a critical component in the treatment of VF or PVT. In adults, delays in time to defibrillation are associated with a relatively linear increase in the risk of mortality of 7% to 10% per minute.<sup>43</sup> However, in a 2018 GWTG-R registry study of hospitalized children with VF or PVT IHCA, no association between time to defibrillation and survival was found.<sup>44</sup> The reasons for this finding are likely multifactorial. Most notably, the median time to the first defibrillation attempt was just 1 minute, which presumably limited the ability to detect timing-driven outcome differences. Moreover, given that 78% of cardiac arrests occurred in the ICU,<sup>44</sup> these children likely received more immediate CPR than adult patients in unmonitored settings. High-quality CPR optimizes myocardial blood flow to the fibrillating myocardium and enhances the likelihood of defibrillation success,<sup>45</sup> so this process likely decreased the implications of modest delays in defibrillation. Nonetheless, being prepared for defibrillation is imperative throughout pediatric CPR events, especially given that many children develop shockable rhythms as a secondary rhythm during CPR.<sup>27</sup>

#### Ventilation and Airway Management

Assisted ventilation is a key component of pediatric CPR, but resuscitation guidelines recommend relatively low ventilation rates during CPR for several reasons.<sup>9,10</sup> First, because of the lower cardiac output and pulmonary blood flow during CPR, less minute ventilation is necessary to balance ventilation and perfusion.<sup>46,47</sup> Second, the delivery of positive-pressure breaths can have deleterious hemodynamic results, most directly through the reduction of systemic venous return.<sup>48</sup> Third, in patients without an invasive airway in place, interruptions in CPR to deliver rescue breaths decrease the chest compression fraction and therefore compromise organ perfusion.<sup>49</sup>

Although based on sound physiological principles, ventilation rates for children that match those recommended for adults do not

take into account that children have higher ventilation rates at baseline or that most pediatric IHCA occur in the setting of respiratory failure.<sup>23,24</sup> A recent observational cohort study of children with IHCA in the ICU found that higher ventilation rates during CPR were associated with improved outcomes.<sup>50</sup> None of the 52 CPR events had AHA guideline-compliant ventilation rates; the median ventilation rate delivered to these children was 30.1 breaths per minute.<sup>50</sup> This finding represents a substantial gap between consensus guidelines and clinical practice, highlighting the need for prospective studies.

Similarly, a clear consensus on optimal pediatric airway management during CPR is needed. As many as 82% of hospitalized children have an invasive airway in place at the time of CPR initiation.<sup>23</sup> A large retrospective study of children without an advanced airway in place during IHCA found an association between the performance of tracheal intubation during cardiac arrest and lower rates of survival.<sup>51</sup> These findings are similar to those of a pediatric OHCA study that demonstrated worse outcomes with endotracheal intubation than with bag-mask ventilation.<sup>52</sup> This association of intubation with worse outcomes may be attributed to interruptions in CPR to facilitate intubation, diverted focus from maintaining CPR quality, unrecognized esophageal intubation and other intubation-associated adverse events, or other unmeasured confounders. When assessing the risk of intubation, pediatric clinicians must consider patient-specific factors as well as their setting and clinical capabilities. The cause of the cardiac arrest and the likelihood of achieving ROSC without invasive ventilation should be considered along with the detrimental implications of interrupting compressions to deliver ventilations.<sup>49</sup> Given the limitations in the quality of data available, the authors of the 2019 AHA guidelines were unable "to make any recommendation about advanced airway management for IHCA."<sup>53</sup>(p907)

### Medications

Epinephrine is recommended during pediatric IHCA because of its  $\alpha$ -adrenergic function of increasing systemic vascular resistance, which increases coronary and cerebral perfusion pressures.<sup>9</sup> No prospective pediatric trial has established the efficacy of epinephrine in improving survival outcomes, but delays in epinephrine administration are associated with worse outcomes in children.<sup>54</sup> Observational findings are mixed regarding the appropriate dosing intervals. The recommended dosing interval of 3 to 5 minutes was associated with improved outcomes in one study,<sup>55</sup> whereas more frequent<sup>56</sup> and less frequent<sup>57</sup> intervals have each been independently associated with superior outcomes in other studies.

Vasopressin, which was previously recommended as an alternative vasopressor for adult cardiac arrest, has not been shown to have a gross survival advantage over epinephrine alone in either adult<sup>58</sup> or pediatric<sup>59</sup> studies. Given that vasopressin induces vasoconstriction through the activation of V1a receptors rather than as an adrenergic agonist, vasopressin could hypothetically be advantageous in specific scenarios in which epinephrine is not efficacious or deleterious.<sup>60</sup> For example, in cardiac arrest associated with catecholamine-refractory vasodilatory shock, vasopressin offers an alternative mechanism of increasing systemic vascular resistance.<sup>61</sup> Similarly, in patients with pulmonary hypertension, the potentially harmful increase in pulmonary vascular resistance associated with epinephrine<sup>62</sup> may be avoided by using a vasoactive drug that largely

sparing the pulmonary vasculature from its effects. Regardless, dedicated studies are needed to characterize potential indications for vasopressin in pediatric resuscitation.

Antiarrhythmic agents are provided during pediatric IHCA for defibrillation-refractory VF or PVT. A GWTG-R registry study that compared lidocaine with amiodarone hydrochloride revealed increased rates of ROSC with lidocaine,<sup>63</sup> but a more recent propensity-matched cohort study found no difference in clinical outcomes between these 2 medications.<sup>64</sup> Given the lack of definitive evidence supporting one antiarrhythmic over another, consensus guidelines conclude that use of either lidocaine or amiodarone is reasonable for shock-refractory VF or PVT.<sup>65</sup>

Other medications commonly used during pediatric IHCA include calcium (eg, calcium gluconate, calcium chloride) and sodium bicarbonate. Despite frequent use, routine administration of either of these medications is associated with decreased rates of survival.<sup>66,67</sup> Each is recommended only for specific indications, such as hyperkalemia and sodium channel overdose.<sup>9</sup>

### Personalizing CPR

Resuscitation techniques and standards have been developed primarily for adult OHCA, with algorithms and guidelines recommending a uniform approach to account for the wide variation in rescuer experience and knowledge base. However, pediatric IHCA is associated with the progression of many disease processes, the pathophysiological characteristics of which persist during CPR. Recognizing and treating the underlying cause of cardiac arrest is imperative for resuscitation success. The immediate cause of cardiac arrest can generally be attributed to one or more of the classic *Hs and Ts* (eg, hypovolemia, hyperkalemia, and tension pneumothorax) and should be promptly addressed. In addition, common underlying pediatric ICU diagnoses, such as congenital heart disease, septic shock, and pulmonary hypertension, may warrant alternative therapies during resuscitation.<sup>9,61,68</sup>

Given that most pediatric in-hospital CPR is performed by trained teams in highly monitored and well-resourced settings,<sup>15</sup> clinicians frequently have the ability to monitor an individual patient's physiological response to CPR and titrate therapies accordingly. The most well-established physiological monitors during resuscitation are invasively measured blood pressure and end-tidal carbon dioxide (EtCO<sub>2</sub>).<sup>9,69</sup> A growing body of literature recognizes that measurement of these physiological parameters can be used to gauge CPR quality, identify likely outcomes, and help guide therapies during resuscitation.<sup>69,70</sup>

Coronary perfusion pressure is the driving force for myocardial blood flow, and higher values during CPR are associated with increased likelihood of ROSC in adult IHCA.<sup>71</sup> Because nearly half of the children with IHCA have an arterial catheter in place,<sup>15</sup> hemodynamic monitoring during CPR is often possible. Diastolic blood pressure is recommended as a feasible surrogate for coronary perfusion pressure,<sup>69</sup> and in a prospective observational trial, intra-arrest diastolic blood pressure of 25 mm Hg or higher in infants and 30 mm Hg or higher in older children was associated with increased rates of survival to discharge and survival with favorable neurological outcome.<sup>23</sup> The ideal methods of optimizing hemodynamics during CPR have not been prospectively studied in humans, but in large animal models of pediatric IHCA, titration of vasopressor drugs according to coronary perfusion pressure during CPR has led

Table 3. Post-Cardiac Arrest Syndrome<sup>9,30,53,83-96</sup>

Key component	Clinical manifestations	Monitoring	Therapies
Brain injury	<ul style="list-style-type: none"> <li>• Encephalopathy</li> <li>• Cerebral edema</li> <li>• Seizures</li> <li>• Myoclonus</li> <li>• Coma</li> </ul>	<ul style="list-style-type: none"> <li>• EEG</li> <li>• Clinical neurological examination</li> <li>• Brain CT</li> <li>• Brain MRI</li> <li>• Near-infrared spectroscopy; transcranial Doppler</li> </ul>	<ul style="list-style-type: none"> <li>• Antiepileptic medications</li> <li>• TTM: 32-34 °C or 36-37.5 °C</li> <li>• Sedation</li> <li>• Oxygen saturation 94%-99%</li> </ul>
Myocardial dysfunction	<ul style="list-style-type: none"> <li>• Left and right ventricular systolic and diastolic dysfunction</li> <li>• Hypotension</li> <li>• Arrhythmias</li> <li>• Low cardiac output</li> <li>• Pulmonary edema</li> </ul>	<ul style="list-style-type: none"> <li>• Arterial line</li> <li>• Echocardiogram</li> <li>• Cardiac monitoring</li> </ul>	<ul style="list-style-type: none"> <li>• Inotropic medications</li> <li>• Antiarrhythmic medications</li> <li>• Electrolyte repletion</li> <li>• Mechanical ventilation</li> </ul>
Systemic ischemia-reperfusion response	<ul style="list-style-type: none"> <li>• Hypotension</li> <li>• Vasoplegia</li> <li>• Hypovolemia</li> <li>• Hyperglycemia/hypoglycemia</li> <li>• Adrenal insufficiency</li> <li>• Coagulopathy</li> <li>• Multisystem organ dysfunction</li> <li>• Pyrexia</li> </ul>	<ul style="list-style-type: none"> <li>• Pulse oximetry</li> <li>• Temperature</li> <li>• Urine output</li> <li>• Laboratory studies (pH; lactate; end-organ perfusion; coagulation; oxygenation/ventilation)</li> <li>• Chest radiograph</li> </ul>	<ul style="list-style-type: none"> <li>• Vasoactive medications</li> <li>• Insulin/dextrose</li> <li>• Corticosteroids</li> <li>• Product replacement</li> <li>• TTM/antipyretic medications</li> <li>• Renal replacement therapy</li> </ul>
Persistent precipitating pathophysiological condition	<ul style="list-style-type: none"> <li>• Ongoing underlying disease process (eg, respiratory failure, sepsis, and pulmonary hypertension)</li> </ul>	<ul style="list-style-type: none"> <li>• Observation/monitoring in ICU setting</li> <li>• Ongoing assessment of progression of underlying illnesses</li> </ul>	<ul style="list-style-type: none"> <li>• Specific to individual pathophysiological processes</li> </ul>

Abbreviations: CT, computed tomography; EEG, electroencephalogram; ICU, intensive care unit; MRI, magnetic resonance imaging; TTM, targeted temperature management.

to superior survival outcomes.<sup>72,73</sup> Given the diverse phenotypes of children with IHCA and the known differences in catecholamine responsiveness in humans, vasopressor doses that are titrated to an individual's physiological response may be advantageous.

In the low-flow state of CPR, EtCO<sub>2</sub> largely reflects pulmonary blood flow and is thereby an indirect marker of cardiac output.<sup>46</sup> Particularly low EtCO<sub>2</sub> values during CPR (ie, <10 mm Hg) are associated with mortality in observational adult studies.<sup>74,75</sup> As with hemodynamics, preclinical studies have demonstrated efficacy in targeting EtCO<sub>2</sub> during CPR.<sup>76</sup> A recent observational study in children did not find an association between EtCO<sub>2</sub> and outcomes.<sup>77</sup> Although limited by statistical power, these findings highlight the need to understand potential confounders in the interpretation of EtCO<sub>2</sub> values, including decrements associated with epinephrine administration and elevated values associated with alveolar hypercapnia in the setting of asphyxial cardiac arrest.<sup>62,78</sup>

#### CPR Duration and Extracorporeal Support

Children who require prolonged CPR have worse outcomes than those with brief cardiac arrests.<sup>22</sup> However, a multicenter cohort study demonstrated a 28% rate of survival to discharge among children who received 30 minutes or more of CPR,<sup>5</sup> and a GWTG-R registry publication reported that 16% of children who received 35 minutes or more of CPR survived to hospital discharge.<sup>22</sup> In each of these studies, most survivors had favorable gross neurological outcomes, suggesting that decisions regarding CPR duration and futility are complex and should not be based on CPR duration alone.

These findings are, in part, attributed to the increased use of extracorporeal CPR (ECPR) as a rescue therapy during cardiac arrest. No prospective trial has compared pediatric ECPR with conventional CPR, but a number of observational studies have identified a survival advantage in children, especially those with congenital cardiac disease, who received ECPR compared with children who received conventional CPR.<sup>79-81</sup> Based on these data, the AHA offered a Class 2b recommendation for ECPR to be considered in

children with IHCA and underlying cardiac diagnoses in settings with "existing ECMO [extracorporeal membrane oxygenation] protocols, expertise, and equipment."<sup>53(pe908)</sup> Although ECPR is also being used for patients without heart disease, evidence of its effectiveness in this group is insufficient.<sup>53</sup>

#### Post-Cardiac Arrest Syndrome and Care

Post-cardiac arrest syndrome (PCAS) is a complex cascade that begins immediately after ROSC and continues for days to weeks. The 4 key components are brain injury, myocardial dysfunction, systemic ischemia-reperfusion injury, and persistent precipitating pathophysiological condition.<sup>30,82</sup>

Clinical manifestations of PCAS range in severity and are summarized in Table 3.<sup>30</sup> Hypoxic-ischemic brain injury can present as encephalopathy or seizures, progress in the days after ROSC, and result in long-term neurodevelopmental deficits in survivors.<sup>30,97</sup> Myocardial dysfunction occurs early after ROSC and, in many patients, peaks at approximately 24 hours before normalizing within 48 to 72 hours after ROSC.<sup>98</sup> Not surprisingly, children with more severe left ventricular dysfunction and those who require a higher vasoactive infusion dose have lower rates of survival to discharge.<sup>83,84</sup> The systemic ischemia-reperfusion response resembles a sepsislike syndrome and can exacerbate myocardial dysfunction and brain injury.<sup>98</sup> The combination of these 3 key components is superimposed on the initial precipitating pathophysiological condition that may persist after ROSC. Identifying and treating the underlying disease process associated with the cardiac arrest can reduce the risk of rearrest.<sup>30,82</sup> Understanding the pathophysiological process allows the clinician to initiate prompt post-cardiac arrest care with the goal of minimizing secondary injury.

#### Therapeutics

Prompt initiation of treatment after IHCA is important to prevent secondary multisystem organ injury (Table 3).<sup>9,30,53,83-96</sup> Initial treatment should focus on normalizing systemic derangements such as

hypotension and hypoxemia. Although patients may be hypertensive immediately after cardiac arrest, hypotension occurs in more than half of children in the first 6 hours after ROSC.<sup>85</sup> At least 1 blood pressure measure below the 5th percentile for age in the first 12 hours after ROSC and longer durations of hypotension during the first 24 hours after ROSC are associated with lower rates of survival to discharge.<sup>86,87</sup> Despite this finding, at least 25% of patients with hypotension do not receive vasoactive infusions.<sup>85</sup> Fluids, inotropes, and vasoactive medications should be administered with the goal of avoiding hypotension by treating cardiac dysfunction and vasoplegia.<sup>9,30</sup>

Post-cardiac arrest hypoxemia is commonly attributed to precipitating lung disease, aspiration, or pulmonary edema and should be anticipated and immediately addressed.<sup>30</sup> Research findings regarding the implications of post-cardiac arrest hyperoxia for outcomes are mixed,<sup>88-90</sup> but it is recommended that clinicians avoid hyperoxia and target normoxia with a goal oxygen saturation of 94% to 99%.<sup>9,30</sup> Studies are similarly in conflict regarding the clinical significance of hypercapnia or hypocapnia after cardiac arrest.<sup>88,91</sup> Guidelines advocate targeting normocapnia or "a PaCO<sub>2</sub> specific for the patient's condition" in the post-ROSC period.<sup>30(p203)</sup> This recommendation is based, in part, on the physiological rationale that hypercapnia may result in cerebral vasodilation and hyperemia and that hypocapnia may cause cerebral vasoconstriction and ischemia.<sup>30,92</sup>

Fever is common after cardiac arrest and is associated with worse outcomes through increasing metabolic demand, promoting free radical production, and other detrimental consequences.<sup>93</sup> Targeted temperature management (TTM) using active temperature regulation devices and antipyretics should be initiated early after ROSC, but therapeutic hypothermia has not been found to offer an advantage over normothermia.<sup>94</sup> The AHA guidelines recommend targeting a temperature of either 32 to 34 °C or 36 to 37.5 °C for 48 hours, followed by 36 to 37.5 °C for an additional 3 days.<sup>30,53</sup>

Seizures are common after cardiac arrest and are often nonconvulsive.<sup>95</sup> Electrographic status epilepticus is associated with worse outcomes.<sup>96</sup> Therefore, continuous electroencephalogram monitoring is recommended for detection and guiding therapy.<sup>30</sup> Antiseizure medications are routinely used to treat seizures and status epilepticus, but it is unclear whether treatment is associated with improved outcomes.

Although most care is supportive, clinicians should consider antibiotics for infection, blood products for coagulopathy-induced bleeding, and corticosteroids for adrenal insufficiency. Close monitoring for organ dysfunction such as acute kidney injury is important to minimize the toxic effects of therapies as patients recover from PCAS.<sup>30</sup> Active areas of investigation for post-cardiac arrest care include advanced neuroimaging, continuous optical measure-

ments of cerebral blood flow and oxygenation, and peripheral biosignatures of neurological injury.<sup>99</sup>

### Identification of Risk and Probable Outcomes

After ROSC, only half of patients survive to discharge, and many survivors are left with substantial neurological morbidities.<sup>3,31</sup> Accurate and timely identification of at-risk patients and probable outcomes enables families and clinicians to make informed decisions regarding goals of care. However, this identification should not be done too early because it can result in incorrect conclusions, and post-cardiac arrest care may lead to improved outcomes.<sup>30</sup> The optimal timing for identification of risk and probable outcomes in children has not been established,<sup>30</sup> but adult guidelines recommend waiting for 72 hours after ROSC in patients who were not treated with TTM and 72 hours after return to normothermia in patients who were treated with TTM.<sup>100</sup> In addition, the neurological examination can be confounded by sedative medications often used during post-cardiac arrest care, and medication clearance may be delayed in the setting of hepatic injury, kidney injury, or therapeutic hypothermia.<sup>101</sup>

Post-cardiac arrest neuroprognostication requires the integration of a patient's pre-cardiac arrest functional status and comorbid medical conditions; IHCA characteristics; and post-cardiac arrest neurological examination, cerebral physiological functions, neuroimaging, and electroencephalogram. Many of these factors have been associated with outcomes, but no single factor has had sufficient accuracy in early identification of risks and probable outcomes after ROSC.<sup>9,30</sup> Serum biomarkers of hypoxic-ischemic brain injury have demonstrated an association with neurological outcomes,<sup>102-104</sup> although further prospective study at serial time points after cardiac arrest is needed before these biomarkers can be incorporated into prognostic algorithms. Patients with catastrophic, irreversible brain injury who do not demonstrate neurological recovery may be evaluated for death by neurological criteria. To undergo such evaluation, patients must be normothermic, have been observed for sufficient time for clearance of any sedative medications, and have no confounding conditions.<sup>105</sup>

## Conclusions

Despite improvements in care and survival in the United States, more than half of children with IHCA die before hospital discharge. Recognition and timely identification of at-risk patients as well as delivery of high-quality CPR and post-cardiac arrest care are imperative to maximize the chances of achieving favorable outcomes. Substantial knowledge gaps remain in how to best care for these vulnerable patients, highlighting the ongoing need for research in this field.

### ARTICLE INFORMATION

**Accepted for Publication:** June 1, 2020.

**Published Online:** November 23, 2020.  
doi:10.1001/jamapediatrics.2020.5039

**Author Contributions:** Drs Morgan and Topjian had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

*Concept and design:* All authors.

*Acquisition, analysis, or interpretation of data:* Morgan, Kirschen, Sutton, Topjian.

*Drafting of the manuscript:* All authors.

*Critical revision of the manuscript for important intellectual content:* All authors.

*Administrative, technical, or material support:* Morgan.

*Supervision:* Morgan, Kilbaugh, Sutton, Topjian.

**Conflict of Interest Disclosures:** Dr Morgan reported receiving grants from the National Institutes of Health (NIH) National Heart, Lung, and Blood Institute (NHLBI) in support of his effort on the submitted work and grants from the NIH NHLBI and Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) outside the submitted work. Dr Sutton reported receiving grants from the NIH NHLBI and grants

from the NICHD outside the submitted work; being chair of the Pediatric Research Task Force of the American Heart Association (AHA) Get With The Guidelines—Resuscitation registry; and being a main author of Pediatric Advanced Life Support Guidelines. Dr Topjian reported receiving grants from the NIH during the conduct of the study, being a volunteer for the AHA, and being chair of the AHA pediatric guidelines writing group. No other disclosures were reported.

**Funding/Support:** This study was supported in part by Career Development Award K23HL148541 from the NIH NHLBI (PI: Dr Morgan).

**Role of the Funder/Sponsor:** The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

## REFERENCES

- Holmberg MJ, Ross CE, Fitzmaurice GM, et al; American Heart Association's Get With The Guidelines—Resuscitation Investigators. Annual incidence of adult and pediatric in-hospital cardiac arrest in the United States. *Circ Cardiovasc Qual Outcomes*. 2019;12(7):e005580. doi:10.1161/CIRCOUTCOMES.119.005580
- Girotra S, Spertus JA, Li Y, Berg RA, Nadkarni VM, Chan PS; American Heart Association Get With The Guidelines—Resuscitation Investigators. Survival trends in pediatric in-hospital cardiac arrests: an analysis from Get With The Guidelines—Resuscitation. *Circ Cardiovasc Qual Outcomes*. 2013;6(1):42-49. doi:10.1161/CIRCOUTCOMES.112.967968
- Holmberg MJ, Wiberg S, Ross CE, et al. Trends in survival after pediatric in-hospital cardiac arrest in the United States. *Circulation*. 2019;140(17):1398-1408. doi:10.1161/CIRCULATIONAHA.119.041667
- Nadkarni VM, Larkin GL, Peberdy MA, et al; National Registry of Cardiopulmonary Resuscitation Investigators. First documented rhythm and clinical outcome from in-hospital cardiac arrest among children and adults. *JAMA*. 2006;295(1):50-57. doi:10.1001/jama.295.1.50
- Berg RA, Nadkarni VM, Clark AE, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Collaborative Pediatric Critical Care Research Network. Incidence and outcomes of cardiopulmonary resuscitation in PICUs. *Crit Care Med*. 2016;44(4):798-808. doi:10.1097/CCM.0000000000001484
- Andersen LW, Holmberg MJ, Berg KM, Donnino MW, Granfeldt A. In-hospital cardiac arrest: a review. *JAMA*. 2019;321(12):1200-1210. doi:10.1001/jama.2019.1696
- Virani SS, Alonso A, Benjamin EJ, et al; American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2020 update: a report from the American Heart Association. *Circulation*. 2020;141(9):e139-e596. doi:10.1161/CIR.0000000000000757
- Fiser DH. Assessing the outcome of pediatric intensive care. *J Pediatr*. 1992;121(1):68-74. doi:10.1016/S0022-3476(05)82544-2
- de Caen AR, Berg MD, Chameides L, et al. Part 12: pediatric advanced life support: 2015 American Heart Association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2015;132(18 suppl 2):S526-S542. doi:10.1161/CIR.0000000000000266
- Atkins DL, Berger S, Duff JP, et al. Part 11: pediatric basic life support and cardiopulmonary resuscitation quality: 2015 American Heart Association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2015;132(18 suppl 2):S519-S525. doi:10.1161/CIR.0000000000000265
- Meert KL, Telford R, Holubkov R, et al; Therapeutic Hypothermia after Pediatric Cardiac Arrest (THAPCA) Trial Investigators. Pediatric out-of-hospital cardiac arrest characteristics and their association with survival and neurobehavioral outcome. *Pediatr Crit Care Med*. 2016;17(12):e543-e550. doi:10.1097/PCC.0000000000000969
- Fink EL, Prince DK, Kaltman JR, et al; Resuscitation Outcomes Consortium. Unchanged pediatric out-of-hospital cardiac arrest incidence and survival rates with regional variation in North America. *Resuscitation*. 2016;107:121-128. doi:10.1016/j.resuscitation.2016.07.244
- McCarthy JJ, Carr B, Sasson C, et al; American Heart Association Emergency Cardiovascular Care Committee; Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation; and the Mission: Lifeline Resuscitation Subcommittee. Out-of-hospital cardiac arrest resuscitation systems of care: a scientific statement from the American Heart Association. *Circulation*. 2018;137(21):e645-e660. doi:10.1161/CIR.0000000000000557
- Nolan JP, Berg RA, Andersen LW, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update of the Utstein resuscitation registry template for in-hospital cardiac arrest: a consensus report from a Task Force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian and New Zealand Council on Resuscitation, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Southern Africa, Resuscitation Council of Asia). *Circulation*. 2019;140(18):e746-e757. doi:10.1161/CIR.0000000000000710
- Berg RA, Sutton RM, Holubkov R, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Collaborative Pediatric Critical Care Research Network and for the American Heart Association's Get With The Guidelines—Resuscitation (formerly the National Registry of Cardiopulmonary Resuscitation) Investigators. Ratio of PICU versus ward cardiopulmonary resuscitation events is increasing. *Crit Care Med*. 2013;41(10):2292-2297. doi:10.1097/CCM.0b013e31828cf0c0
- Tibballs J, Kinney S. Reduction of hospital mortality and of preventable cardiac arrest and death on introduction of a pediatric medical emergency team. *Pediatr Crit Care Med*. 2009;10(3):306-312. doi:10.1097/PCC.0b013e318198b02c
- Sen AI, Morgan RW, Morris MC. Variability in the implementation of rapid response teams at academic American pediatric hospitals. *J Pediatr*. 2013;163(6):1772-1774. doi:10.1016/j.jpeds.2013.07.018
- Brilli RJ, Gibson R, Luria JW, et al. Implementation of a medical emergency team in a large pediatric teaching hospital prevents respiratory and cardiopulmonary arrests outside the intensive care unit. *Pediatr Crit Care Med*. 2007;8(3):236-246. doi:10.1097/01.PCC.00000262947.72442.EA
- Chapman SM, Wray J, Oulton K, Peters MJ. Systematic review of paediatric track and trigger systems for hospitalised children. *Resuscitation*. 2016;109:87-109. doi:10.1016/j.resuscitation.2016.07.230
- Knudson JD, Neish SR, Cabrera AG, et al. Prevalence and outcomes of pediatric in-hospital cardiopulmonary resuscitation in the United States: an analysis of the Kids' Inpatient Database\*. *Crit Care Med*. 2012;40(11):2940-2944. doi:10.1097/CCM.0b013e31825feb3f
- Alten JA, Klugman D, Raymond TT, et al. Epidemiology and outcomes of cardiac arrest in pediatric cardiac ICUs. *Pediatr Crit Care Med*. 2017;18(10):935-943. doi:10.1097/PCC.0000000000001273
- Matos RI, Watson RS, Nadkarni VM, et al; American Heart Association's Get With The Guidelines—Resuscitation (Formerly the National Registry of Cardiopulmonary Resuscitation) Investigators. Duration of cardiopulmonary resuscitation and illness category impact survival and neurologic outcomes for in-hospital pediatric cardiac arrests. *Circulation*. 2013;127(4):442-451. doi:10.1161/CIRCULATIONAHA.112.125625
- Berg RA, Sutton RM, Reeder RW, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Collaborative Pediatric Critical Care Research Network (CPCCRN) PICUcPR (Pediatric Intensive Care Quality of Cardio-Pulmonary Resuscitation) Investigators. Association between diastolic blood pressure during pediatric in-hospital cardiopulmonary resuscitation and survival. *Circulation*. 2018;137(17):1784-1795. doi:10.1161/CIRCULATIONAHA.117.032270
- Khera R, Tang Y, Girotra S, et al; American Heart Association's Get With The Guidelines—Resuscitation Investigators. Pulselessness after initiation of cardiopulmonary resuscitation for bradycardia in hospitalized children. *Circulation*. 2019;140(5):370-378. doi:10.1161/CIRCULATIONAHA.118.039048
- Donoghue A, Berg RA, Hazinski MF, Praestgaard AH, Roberts K, Nadkarni VM; American Heart Association National Registry of CPR Investigators. Cardiopulmonary resuscitation for bradycardia with poor perfusion versus pulseless cardiac arrest. *Pediatrics*. 2009;124(6):1541-1548. doi:10.1542/peds.2009-0727
- Morgan RW, Reeder RW, Meert KL, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development/Collaborative Pediatric Critical Care Research Network (CPCCRN) Pediatric Intensive Care Quality of Cardio-Pulmonary Resuscitation (PICUcPR) Investigators. Survival and hemodynamics during pediatric cardiopulmonary resuscitation for bradycardia and poor perfusion versus pulseless cardiac arrest. *Crit Care Med*. 2020;48(6):881-889. doi:10.1097/CCM.0000000000004308
- Samson RA, Nadkarni VM, Meaney PA, Carey SM, Berg MD, Berg RA; American Heart Association National Registry of CPR Investigators. Outcomes

- of in-hospital ventricular fibrillation in children. *N Engl J Med*. 2006;354(22):2328-2339. doi:10.1056/NEJMoa052917
28. Sutton RM, French B, Niles DE, et al. 2010 American Heart Association recommended compression depths during pediatric in-hospital resuscitations are associated with survival. *Resuscitation*. 2014;85(9):1179-1184. doi:10.1016/j.resuscitation.2014.05.007
29. Sutton RM, French B, Nishisaki A, et al. American Heart Association cardiopulmonary resuscitation quality targets are associated with improved arterial blood pressure during pediatric cardiac arrest. *Resuscitation*. 2013;84(2):168-172. doi:10.1016/j.resuscitation.2012.08.335
30. Topjian AA, de Caen A, Wainwright MS, et al. Pediatric post-cardiac arrest care: a scientific statement from the American Heart Association. *Circulation*. 2019;140(6):e194-e233. doi:10.1161/CIR.0000000000000697
31. Slomine BS, Silverstein FS, Christensen JR, et al; Therapeutic Hypothermia after Paediatric Cardiac Arrest (THAPCA) Trial Investigators. Neurobehavioural outcomes in children after in-hospital cardiac arrest. *Resuscitation*. 2018;124:80-89. doi:10.1016/j.resuscitation.2018.01.002
32. Du Pont-Thibodeau G, Fry M, Kirschen M, et al. Timing and modes of death after pediatric out-of-hospital cardiac arrest resuscitation. *Resuscitation*. 2018;133:160-166. doi:10.1016/j.resuscitation.2018.08.014
33. Brady PW, Muething S, Kotagal U, et al. Improving situation awareness to reduce unrecognized clinical deterioration and serious safety events. *Pediatrics*. 2013;131(1):e298-e308. doi:10.1542/peds.2012-1364
34. Bonafide CP, Roberts KE, Priestley MA, et al. Development of a pragmatic measure for evaluating and optimizing rapid response systems. *Pediatrics*. 2012;129(4):e874-e881. doi:10.1542/peds.2011-2784
35. Parshuram CS, Dryden-Palmer K, Farrell C, et al; Canadian Critical Care Trials Group and the EPOCH Investigators. Effect of a pediatric early warning system on all-cause mortality in hospitalized pediatric patients: the EPOCH randomized clinical trial. *JAMA*. 2018;319(10):1002-1012. doi:10.1001/jama.2018.0948
36. Dewan M, Muthu N, Shelov E, et al. Performance of a clinical decision support tool to identify PICU patients at high risk for clinical deterioration. *Pediatr Crit Care Med*. 2020;21(2):129-135. doi:10.1097/PCC.0000000000002106
37. Niles DE, Dewan M, Zebuhr C, et al. A pragmatic checklist to identify pediatric ICU patients at risk for cardiac arrest or code bell activation. *Resuscitation*. 2016;99:33-37. doi:10.1016/j.resuscitation.2015.11.017
38. Alten J, Klugman D, Cooper DS, et al. Abstract 11798: cardiac arrest prevention quality improvement project from the pediatric cardiac critical care consortium. *Circulation*. 2019;140:A11798.
39. Gupta P, Rettiganti M, Rice TB, Wetzel RC. Impact of 24/7 in-hospital intensivist coverage on outcomes in pediatric intensive care: a multicenter study. *Am J Respir Crit Care Med*. 2016;194(12):1506-1513. doi:10.1164/rccm.201512-2456OC
40. Wolfe H, Zebuhr C, Topjian AA, et al. Interdisciplinary ICU cardiac arrest debriefing improves survival outcomes\*. *Crit Care Med*. 2014;42(7):1688-1695. doi:10.1097/CCM.0000000000000327
41. Sutton RM, Reeder RW, Landis W, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Collaborative Pediatric Critical Care Research Network (CPCCRN) Investigators. Chest compression rates and pediatric in-hospital cardiac arrest survival outcomes. *Resuscitation*. 2018;130:159-166. doi:10.1016/j.resuscitation.2018.07.015
42. Niles DE, Duval-Arnould J, Skellet S, et al; Pediatric Resuscitation Quality (pediRES-Q) Collaborative Investigators. Characterization of pediatric in-hospital cardiopulmonary resuscitation quality metrics across an international resuscitation collaborative. *Pediatr Crit Care Med*. 2018;19(5):421-432. doi:10.1097/PCC.0000000000001520
43. Chan PS, Krumholz HM, Nichol G, Nallamothu BK; American Heart Association National Registry of Cardiopulmonary Resuscitation Investigators. Delayed time to defibrillation after in-hospital cardiac arrest. *N Engl J Med*. 2008;358(1):9-17. doi:10.1056/NEJMoa0706467
44. Hunt EA, Duval-Arnould JM, Bembea MM, et al; American Heart Association's Get With The Guidelines-Resuscitation Investigators. Association between time to defibrillation and survival in pediatric in-hospital cardiac arrest with a first documented shockable rhythm. *JAMA Netw Open*. 2018;1(5):e182643. doi:10.1001/jamanetworkopen.2018.2643
45. Berg RA, Hilwig RW, Berg MD, et al. Immediate post-shock chest compressions improve outcome from prolonged ventricular fibrillation. *Resuscitation*. 2008;78(1):71-76. doi:10.1016/j.resuscitation.2008.02.014
46. Weil MH, Bisera J, Trevino RP, Rackow EC. Cardiac output and end-tidal carbon dioxide. *Crit Care Med*. 1985;13(11):907-909. doi:10.1097/00003246-198511000-00011
47. Maier GW, Newton JR Jr, Wolfe JA, et al. The influence of manual chest compression rate on hemodynamic support during cardiac arrest: high-impulse cardiopulmonary resuscitation. *Circulation*. 1986;74(6 Pt 2):IV51-IV59.
48. Aufderheide TP, Sigurdsson G, Pirrallo RG, et al. Hyperventilation-induced hypotension during cardiopulmonary resuscitation. *Circulation*. 2004;109(16):1960-1965. doi:10.1161/01.CIR.0000126594.79136.61
49. Berg RA, Sanders AB, Kern KB, et al. Adverse hemodynamic effects of interrupting chest compressions for rescue breathing during cardiopulmonary resuscitation for ventricular fibrillation cardiac arrest. *Circulation*. 2001;104(20):2465-2470. doi:10.1161/hc4501.098926
50. Sutton RM, Reeder RW, Landis WP, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Collaborative Pediatric Critical Care Research Network (CPCCRN). Ventilation rates and pediatric in-hospital cardiac arrest survival outcomes. *Crit Care Med*. 2019;47(11):1627-1636. doi:10.1097/CCM.0000000000003898
51. Andersen LW, Raymond TT, Berg RA, et al; American Heart Association's Get With The Guidelines-Resuscitation Investigators. Association between tracheal intubation during pediatric in-hospital cardiac arrest and survival. *JAMA*. 2016;316(17):1786-1797. doi:10.1001/jama.2016.14486
52. Hansen ML, Lin A, Eriksson C, et al; CARES surveillance group. A comparison of pediatric airway management techniques during out-of-hospital cardiac arrest using the CARES database. *Resuscitation*. 2017;120:51-56. doi:10.1016/j.resuscitation.2017.08.015
53. Duff JP, Topjian AA, Berg MD, et al. 2019 American Heart Association focused update on pediatric advanced life support: an update to the American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2019;140(24):e904-e914. doi:10.1161/CIR.0000000000000731
54. Andersen LW, Berg KM, Saindon BZ, et al; American Heart Association Get With The Guidelines-Resuscitation Investigators. Time to epinephrine and survival after pediatric in-hospital cardiac arrest. *JAMA*. 2015;314(8):802-810. doi:10.1001/jama.2015.9678
55. Meert K, Telford R, Holubkov R, et al. Paediatric in-hospital cardiac arrest: factors associated with survival and neurobehavioural outcome one year later. *Resuscitation*. 2018;124:96-105. doi:10.1016/j.resuscitation.2018.01.013
56. Grunau B, Kawano T, Scheuermeyer FX, et al. The association of the average epinephrine dosing interval and survival with favorable neurologic status at hospital discharge in out-of-hospital cardiac arrest. *Ann Emerg Med*. 2019;74(6):797-806. doi:10.1016/j.annemergmed.2019.04.031
57. Hoyme DB, Patel SS, Samson RA, et al; American Heart Association Get With The Guidelines-Resuscitation Investigators. Epinephrine dosing interval and survival outcomes during pediatric in-hospital cardiac arrest. *Resuscitation*. 2017;117:18-23. doi:10.1016/j.resuscitation.2017.05.023
58. Stiell IG, Hébert PC, Wells GA, et al. Vasopressin versus epinephrine for in-hospital cardiac arrest: a randomised controlled trial. *Lancet*. 2001;358(9276):105-109. doi:10.1016/S0140-6736(01)05328-4
59. Duncan JM, Meaney P, Simpson P, Berg RA, Nadkarni V, Schexnayder S; National Registry of CPR Investigators. Vasopressin for in-hospital pediatric cardiac arrest: results from the American Heart Association National Registry of Cardiopulmonary Resuscitation. *Pediatr Crit Care Med*. 2009;10(2):191-195. doi:10.1097/PCC.0b013e31819a36f2
60. Lindner KH, Prenal AW, Pfenninger EG, et al. Vasopressin improves vital organ blood flow during closed-chest cardiopulmonary resuscitation in pigs. *Circulation*. 1995;91(1):215-221. doi:10.1161/01.CIR.91.1.215
61. Morgan RW, Fitzgerald JC, Weiss SL, Nadkarni VM, Sutton RM, Berg RA. Sepsis-associated in-hospital cardiac arrest: epidemiology, pathophysiology, and potential therapies. *J Crit Care*. 2017;40:128-135. doi:10.1016/j.jccr.2017.03.023
62. Lindberg L, Liao Q, Steen S. The effects of epinephrine/norepinephrine on end-tidal carbon dioxide concentration, coronary perfusion pressure and pulmonary arterial blood flow during cardiopulmonary resuscitation. *Resuscitation*. 2000;43(2):129-140. doi:10.1016/S0300-9572(99)00129-X
63. Valdes SO, Donoghue AJ, Hoyme DB, et al; American Heart Association Get With The Guidelines-Resuscitation Investigators. Outcomes



associated with amiodarone and lidocaine in the treatment of in-hospital pediatric cardiac arrest with pulseless ventricular tachycardia or ventricular fibrillation. *Resuscitation*. 2014;85(3):381-386. doi:10.1016/j.resuscitation.2013.12.008

64. Holmberg MJ, Ross CE, Atkins DL, Valdes SO, Donnino MW, Andersen LW; American Heart Association's Get With The Guidelines®-Resuscitation Pediatric Research Task Force. Lidocaine versus amiodarone for pediatric in-hospital cardiac arrest: an observational study. *Resuscitation*. 2020;149:191-201. doi:10.1016/j.resuscitation.2019.12.033

65. McBride ME, Marino BS, Webster G, et al. Amiodarone versus lidocaine for pediatric cardiac arrest due to ventricular arrhythmias: a systematic review. *Pediatr Crit Care Med*. 2017;18(2):183-189. doi:10.1097/PCC.0000000000001026

66. Srinivasan V, Morris MC, Helfaer MA, Berg RA, Nadkarni VM; American Heart Association National Registry of CPR Investigators. Calcium use during in-hospital pediatric cardiopulmonary resuscitation: a report from the National Registry of Cardiopulmonary Resuscitation. *Pediatrics*. 2008; 121(5):e1144-e1151. doi:10.1542/peds.2007-1555

67. Raymond TT, Stromberg D, Stigall W, Burton G, Zaritsky A; American Heart Association's Get With The Guidelines-Resuscitation Investigators. Sodium bicarbonate use during in-hospital pediatric pulseless cardiac arrest—a report from the American Heart Association Get With The Guidelines-Resuscitation. *Resuscitation*. 2015;89: 106-113. doi:10.1016/j.resuscitation.2015.01.007

68. Marino BS, Tabbutt S, MacLaren G, et al; American Heart Association Congenital Cardiac Defects Committee of the Council on Cardiovascular Disease in the Young; Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; Council on Cardiovascular Surgery and Anesthesia; and Emergency Cardiovascular Care Committee. Cardiopulmonary resuscitation in infants and children with cardiac disease: a scientific statement from the American Heart Association. *Circulation*. 2018;137(22):e691-e782. doi:10.1161/CIR.0000000000000524

69. Meaney PA, Bobrow BJ, Mancini ME, et al; CPR Quality Summit Investigators, the American Heart Association Emergency Cardiovascular Care Committee, and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. Cardiopulmonary resuscitation quality: [corrected] improving cardiac resuscitation outcomes both inside and outside the hospital: a consensus statement from the American Heart Association. *Circulation*. 2013;128(4):417-435. doi:10.1161/CIR.0b013e31829d8654

70. Marquez AM, Morgan RW, Ross CE, Berg RA, Sutton RM. Physiology-directed cardiopulmonary resuscitation: advances in precision monitoring during cardiac arrest. *Curr Opin Crit Care*. 2018;24 (3):143-150. doi:10.1097/MCC.0000000000000499

71. Paradis NA, Martin GB, Rivers EP, et al. Coronary perfusion pressure and the return of spontaneous circulation in human cardiopulmonary resuscitation. *JAMA*. 1990;263(8):1106-1113. doi:10.1001/jama.1990.03440080084029

72. Morgan RW, Kilbaugh TJ, Shoap W, et al; Pediatric Cardiac Arrest Survival Outcomes PICASO Laboratory Investigators. A hemodynamic-directed

approach to pediatric cardiopulmonary resuscitation (HD-CPR) improves survival. *Resuscitation*. 2017;111:41-47. doi:10.1016/j.resuscitation.2016.11.018

73. Lautz AJ, Morgan RW, Karlsson M, et al. Hemodynamic-directed cardiopulmonary resuscitation improves neurologic outcomes and mitochondrial function in the heart and brain. *Crit Care Med*. 2019;47(3):e241-e249. doi:10.1097/CCM.00000000000003620

74. Levine RL, Wayne MA, Miller CC. End-tidal carbon dioxide and outcome of out-of-hospital cardiac arrest. *N Engl J Med*. 1997;337(5):301-306. doi:10.1056/NEJM19970713370503

75. Sanders AB, Kern KB, Otto CW, Milander MM, Ewy GA. End-tidal carbon dioxide monitoring during cardiopulmonary resuscitation: a prognostic indicator for survival. *JAMA*. 1989;262(10):1347-1351. doi:10.1001/jama.1989.03430100081033

76. Hamrick JT, Hamrick JL, Bhalala U, et al. End-tidal CO<sub>2</sub>-guided chest compression delivery improves survival in a neonatal asphyxial cardiac arrest model. *Pediatr Crit Care Med*. 2017;18(11): e575-e584. doi:10.1097/PCC.0000000000001299

77. Berg RA, Reeder RW, Meert KL, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Collaborative Pediatric Critical Care Research Network (CPCCRN) Pediatric Intensive Care Quality of Cardio-Pulmonary Resuscitation (PICQPR) investigators. End-tidal carbon dioxide during pediatric in-hospital cardiopulmonary resuscitation. *Resuscitation*. 2018; 133:173-179. doi:10.1016/j.resuscitation.2018.08.013

78. Lah K, Križmarčić M, Grmec S. The dynamic pattern of end-tidal carbon dioxide during cardiopulmonary resuscitation: difference between asphyxial cardiac arrest and ventricular fibrillation/pulseless ventricular tachycardia cardiac arrest. *Crit Care*. 2011;15(1):R13. doi:10.1186/cc9417

79. Ortmann L, Prophan P, Gossett J, et al; American Heart Association's Get With The Guidelines-Resuscitation Investigators. Outcomes after in-hospital cardiac arrest in children with cardiac disease: a report from Get With The Guidelines-Resuscitation. *Circulation*. 2011;124(21): 2329-2337. doi:10.1161/CIRCULATIONAHA.110.013466

80. Lasa JJ, Rogers RS, Localio R, et al. Extracorporeal cardiopulmonary resuscitation (E-CPR) during pediatric in-hospital cardiopulmonary arrest is associated with improved survival to discharge: a report from the American Heart Association's Get With The Guidelines-Resuscitation (GWTG-R) registry. *Circulation*. 2016;133(2):165-176. doi:10.1161/CIRCULATIONAHA.115.016082

81. Bembea MM, Ng DK, Rizkalla N, et al; American Heart Association's Get With The Guidelines – Resuscitation Investigators. Outcomes after extracorporeal cardiopulmonary resuscitation of pediatric in-hospital cardiac arrest: a report from the Get With The Guidelines-Resuscitation and the Extracorporeal Life Support Organization registries. *Crit Care Med*. 2019;47(4):e278-e285. doi:10.1097/CCM.00000000000003622

82. Neumar RW, Nolan JP, Adrie C, et al. Post-cardiac arrest syndrome: epidemiology, pathophysiology, treatment, and prognostication: a consensus statement from the International Liaison Committee on Resuscitation (American Heart Association, Australian and New Zealand

Council on Resuscitation, European Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Asia, and the Resuscitation Council of Southern Africa); the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Perioperative, and Critical Care; the Council on Clinical Cardiology; and the Stroke Council. *Circulation*. 2008;118(23): 2452-2483. doi:10.1161/CIRCULATIONAHA.108.190652

83. Conlon TW, Falkensammer CB, Hammond RS, Nadkarni VM, Berg RA, Topjian AA. Association of left ventricular systolic function and vasopressor support with survival following pediatric out-of-hospital cardiac arrest. *Pediatr Crit Care Med*. 2015;16(2):146-154. doi:10.1097/PCC.0000000000000305

84. Checchia PA, Sehra R, Moynihan J, Daher N, Tang W, Weil MH. Myocardial injury in children following resuscitation after cardiac arrest. *Resuscitation*. 2003;57(2):131-137. doi:10.1016/S0300-9572(03)00003-0

85. Topjian AA, French B, Sutton RM, et al. Early postresuscitation hypotension is associated with increased mortality following pediatric cardiac arrest. *Crit Care Med*. 2014;42(6):1518-1523. doi:10.1097/CCM.0000000000000216

86. Topjian AA, Telford R, Holubkov R, et al; Therapeutic Hypothermia After Pediatric Cardiac Arrest (THAPCA) Trial Investigators. Association of early postresuscitation hypotension with survival to discharge after targeted temperature management for pediatric out-of-hospital cardiac arrest: secondary analysis of a randomized clinical trial. *JAMA Pediatr*. 2018;172(2):143-153. doi:10.1001/jamapediatrics.2017.4043

87. Laverriere EK, Polansky M, French B, Nadkarni VM, Berg RA, Topjian AA. Association of duration of hypotension with survival after pediatric cardiac arrest. *Pediatr Crit Care Med*. 2020;21(2):143-149. doi:10.1097/PCC.0000000000002119

88. Del Castillo J, López-Herce J, Matamoros M, et al; Iberoamerican Pediatric Cardiac Arrest Study Network RIBEPCI. Hyperoxia, hypoxemia and hypercapnia as outcome factors after cardiac arrest in children. *Resuscitation*. 2012;83(12):1456-1461. doi:10.1016/j.resuscitation.2012.07.019

89. Ferguson LP, Durward A, Tibby SM. Relationship between arterial partial oxygen pressure after resuscitation from cardiac arrest and mortality in children. *Circulation*. 2012;126(3):335-342. doi:10.1161/CIRCULATIONAHA.111.085100

90. Holmberg MJ, Nicholson T, Nolan JP, et al; Adult Pediatric Advanced Life Support Task Forces at the International Liaison Committee on Resuscitation (ILCOR). Oxygenation and ventilation targets after cardiac arrest: a systematic review and meta-analysis. *Resuscitation*. 2020;152:107-115. doi:10.1016/j.resuscitation.2020.04.031

91. Bennett KS, Clark AE, Meert KL, et al; Pediatric Emergency Care Medicine Applied Research Network. Early oxygenation and ventilation measurements after pediatric cardiac arrest: lack of association with outcome. *Crit Care Med*. 2013;41 (6):1534-1542. doi:10.1097/CCM.0b013e318287f54c

92. de Caen AR, Maconochie IK, Aickin R, et al; Pediatric Basic Life Support and Pediatric Advanced Life Support Chapter Collaborators. Part 6:

- pediatric basic life support and pediatric advanced life support: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2015;132(suppl 1):S177-S203. doi:10.1161/CIR.0000000000000275
- 93.** Bembea MM, Nadkarni VM, Diener-West M, et al; American Heart Association National Registry of Cardiopulmonary Resuscitation Investigators. Temperature patterns in the early postresuscitation period after pediatric in-hospital cardiac arrest. *Pediatr Crit Care Med*. 2010;11(6):723-730. doi:10.1097/PCC.0b013e3181d6659
- 94.** Moler FW, Silverstein FS, Holubkov R, et al; THAPCA Trial Investigators. Therapeutic hypothermia after in-hospital cardiac arrest in children. *N Engl J Med*. 2017;376(4):318-329. doi:10.1056/NEJMoa1610493
- 95.** Abend NS, Topjian A, Ichord R, et al. Electroencephalographic monitoring during hypothermia after pediatric cardiac arrest. *Neurology*. 2009;72(22):1931-1940. doi:10.1212/WNL.0b013e3181a82687
- 96.** Topjian AA, Sánchez SM, Shults J, Berg RA, Dlugos DJ, Abend NS. Early electroencephalographic background features predict outcomes in children resuscitated from cardiac arrest. *Pediatr Crit Care Med*. 2016;17(6):547-557. doi:10.1097/PCC.0000000000000740
- 97.** Ichord R, Silverstein FS, Slomine BS, et al; THAPCA Trial Group. Neurologic outcomes in pediatric cardiac arrest survivors enrolled in the THAPCA trials. *Neurology*. 2018;91(2):e123-e131. doi:10.1212/WNL.0000000000005773
- 98.** Adrie C, Adib-Conquy M, Laurent I, et al. Successful cardiopulmonary resuscitation after cardiac arrest as a "sepsis-like" syndrome. *Circulation*. 2002;106(5):562-568. doi:10.1161/01.CIR.0000023891.80661.AD
- 99.** Hosseini M, Wilson RH, Crouzet C, Amirhekmat A, Wei KS, Akbari Y. Resuscitating the globally ischemic brain: TTM and beyond. *Neurotherapeutics*. 2020;17(2):539-562. doi:10.1007/s13311-020-00856-z
- 100.** Callaway CW, Donnino MW, Fink EL, et al. Part 8: post-cardiac arrest care: 2015 American Heart Association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2015;132(suppl 2):S465-S482. doi:10.1161/CIR.0000000000000262
- 101.** Link MS, Berkow LC, Kudenchuk PJ, et al. Part 7: adult advanced cardiovascular life support: 2015 American Heart Association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2015;132(suppl 2):S444-S464. doi:10.1161/CIR.0000000000000261
- 102.** Fink EL, Berger RP, Clark RS, et al. Serum biomarkers of brain injury to classify outcome after pediatric cardiac arrest\*. *Crit Care Med*. 2014;42(3):664-674. doi:10.1097/01.ccm.0000435668.53188.80
- 103.** Topjian AA, Lin R, Morris MC, et al. Neuron-specific enolase and S-100B are associated with neurologic outcome after pediatric cardiac arrest. *Pediatr Crit Care Med*. 2009;10(4):479-490. doi:10.1097/PCC.0b013e318198bdb5
- 104.** Kirschen MP, Yehya N, Graham K, et al. Circulating neurofilament light chain is associated with survival after pediatric cardiac arrest. *Pediatr Crit Care Med*. 2020;21(7):656-661. doi:10.1097/PCC.0000000000002294
- 105.** Nakagawa TA, Ashwal S, Mathur M, et al; Society of Critical Care Medicine; Section on Critical Care and Section on Neurology of the American Academy of Pediatrics; Child Neurology Society. Guidelines for the determination of brain death in infants and children: an update of the 1987 Task Force recommendations. *Crit Care Med*. 2011;39(9):2139-2155. doi:10.1097/CCM.0b013e31821f0d4f