

Letters

RESEARCH LETTER

Psychological Distress and Loneliness Reported by US Adults in 2018 and April 2020

Coronavirus disease 2019 (COVID-19) introduced stressors to mental health, including loneliness stemming from social isolation, fear of contracting the disease, economic strain, and uncertainty about the future. We fielded a national survey measuring symptoms of psychological distress and loneliness among US adults in April 2020 and compared results with national data from 2018.

Methods | We fielded the Johns Hopkins COVID-19 Civic Life and Public Health Survey from April 7 to April 13, 2020, using NORC's AmeriSpeak Panel. AmeriSpeak is a probability-based panel designed to be representative of the US adult population. The panel is sourced from NORC's area probability sample and from a US Postal Service address-based sample covering 97% of US households.¹ The panel has a recruitment rate of 34% and includes approximately 35 000 members. The sample for the Johns Hopkins survey was drawn from this panel and the survey was administered online. NORC obtains informed consent prior to enrolling individuals in the panel. The Johns Hopkins Bloomberg School of Public Health institutional review board deemed this study not human participants research and waived informed consent.

We measured the prevalence of symptoms of serious psychological distress in the overall sample and among demographic subgroups using the Kessler 6 Psychological Distress Scale, with the validated measure of serious distress defined as a score of 13 or higher on the 0- to 24-point scale.² We also measured the proportion of respondents who reported that they always or often feel lonely in response to the item "How often do you feel lonely?" with response options always, often, sometimes, rarely, and never.


We compared the prevalence of symptoms of serious psychological distress in April 2020 with an identical measure from the 2018 National Health Interview Survey (NHIS), which used the Kessler 6 scale among 25 417 adults aged 18 years or older in household interviews. The 2018 NHIS response rate was 64.2%.³

For each measure, we calculated proportions and 95% CIs using Stata version 15 (StataCorp). The Johns Hopkins and NHIS survey data were analyzed separately. Analyses of both data sets incorporated survey sampling weights to generate nationally representative estimates.

Results | The survey response rate was 70.4%, with a final sample of 1468 adults aged 18 years or older.

In April 2020, 13.6% (95% CI, 11.1%-16.5%) of US adults reported symptoms of serious psychological distress, relative to 3.9% (95% CI, 3.6%-4.2%) in 2018 (Figure). Among the subgroups examined, in April 2020, symptoms of psychological

Figure. Psychological Distress Among US Adults Aged 18 Years or Older Overall and by Subgroup, April 2020 vs 2018



April 2020 measures are from wave 1 of the Johns Hopkins COVID-19 Civic Life and Public Health Survey, fielded April 7-13, 2020 (N = 1468 adults aged ≥18 years). 2018 Measures of psychological distress are from the 2018 National Health Interview Survey (N = 25 417 adults aged ≥18 years). Psychological distress was measured using the Kessler 6 Psychological Distress Scale, with scores of 13 or higher indicating serious psychological distress. The error bars indicate 95% CIs.

^a Race/ethnicity was collected as part of the demographic profile in both the April 2020 Johns Hopkins survey and the 2018 National Health Interview Survey. In both surveys, the options were defined by the study investigators, and participants classified their own race/ethnicity.

distress were highest among young adults aged 18 to 29 years (24.0% [95% CI, 16.3%-33.8%]), adults with household income of less than \$35 000 per year (19.3% [95% CI, 14.2%-25.6%]), and Hispanic adults (18.3% [95% CI, 11.2%-28.3%]).

The corresponding prevalence estimates for these 3 groups in 2018 were 3.7% (95% CI, 3.0%-4.6%), 7.9% (95% CI, 7.1%-8.6%), and 4.4% (95% CI, 3.7%-5.4%), respectively. The lowest prevalence of serious psychological distress among the subgroups examined in April 2020 was observed in adults aged 55 years or older (7.3% [95% CI, 4.8%-10.9%]). In April 2020, 13.8% (95% CI, 11.4%-16.6%) of US adults reported that they always or often felt lonely.

Discussion | The prevalence of reported symptoms of psychological distress among US adults was higher in 2020 during the COVID-19 pandemic than in 2018. This finding builds on prior research documenting psychological distress among health care workers responding to COVID-19.⁴

The measure of serious psychological distress derived from the Kessler 6 scale has been shown to accurately predict serious mental illness,² suggesting acute distress during COVID-19 may transfer to longer-term psychiatric disorders. In April 2020, 13.8% of US adults reported that they always or often felt lonely. In comparison, a national survey using an identical measure of loneliness found that 11% of US adults reported always or often feeling lonely in April and May 2018.⁵ Because loneliness increased only slightly from 2018 to 2020, other factors may be driving psychological distress during the COVID-19 pandemic.

The NORC AmeriSpeak panel used probability-based recruitment consistent with best-practice standards for survey research,⁶ but results may be vulnerable to sampling biases. The degree to which US adults classified as essential workers during the COVID-19 pandemic were represented in the survey sample is unknown. While both surveys are designed to be nationally representative of US adults, the sampling and recruitment methods and mode of administration varied in the Johns Hopkins April 2020 and NHIS 2018 surveys. There is a potential for selection bias if individuals were more likely to respond to a survey about psychological distress in April 2020 vs 2018.

Emma E. McGinty, PhD
Rachel Presskreischer, MS
Hahrie Han, PhD
Colleen L. Barry, PhD

Author Affiliations: Department of Health Policy and Management, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland (McGinty, Presskreischer, Barry); Department of Political Science, Johns Hopkins University, Baltimore, Maryland (Han).

Corresponding Author: Emma E. McGinty, PhD, Department of Health Policy and Management, Johns Hopkins Bloomberg School of Public Health, 624 N Broadway, Room 359, Baltimore, MD 21205 (bmcginty@jhu.edu).

Accepted for Publication: May 19, 2020.

Published Online: June 3, 2020. doi:10.1001/jama.2020.9740

Author Contributions: Dr McGinty had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: McGinty, Han, Barry.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: McGinty.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: McGinty, Presskreischer.

Obtained funding: McGinty, Han, Barry.

Administrative, technical, or material support: Barry.

Supervision: McGinty, Han, Barry.

Conflict of Interest Disclosures: None reported.

Funding/Support: Dr McGinty reported receiving a Faculty Innovation Award from Johns Hopkins University. Dr Han reported receiving a grant from the Robert Wood Johnson Foundation. Dr Barry reported receiving endowment funds from the Johns Hopkins Bloomberg School of Public Health.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

1. Dennis JM. *Technical Overview of the AmeriSpeak Panel NORC's Probability-Based Household Panel*. NORC at the University of Chicago; 2019.
2. Kessler RC, Barker PR, Colpe LJ, et al. Screening for serious mental illness in the general population. *Arch Gen Psychiatry*. 2003;60(2):184-189. doi:10.1001/archpsyc.60.2.184
3. Centers for Disease Control and Prevention. National Center for Health Statistics: data, questionnaires, and related documentation. Accessed May 13, 2020. <https://www.cdc.gov/nchs/nhis/data-questionnaires-documentation.htm>
4. Lai J, Ma S, Wang Y, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Network Open*. 2020;3(3):e203976-e.
5. Kaiser Family Foundation; The Economist. Survey on Loneliness and Social Isolation in the United States, the United Kingdom, and Japan. Published 2018. Accessed May 13, 2020. <http://files.kff.org/attachment/Topline-Kaiser-Family-Foundation-The-Economist-Survey-on-Loneliness-and-Social-Isolation-in-the-United-States-the-United-Kingdom-and-Japan>
6. American Association for Public Opinion Research. Report on online panels. Published June 2010. Accessed May 14, 2020. <https://www.aapor.org/Education-Resources/Reports/Report-on-Online-Panels.aspx>

Correlation Between N95 Extended Use and Reuse and Fit Failure in an Emergency Department

Frontline health care workers are at high risk of contracting severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which causes coronavirus disease 2019 (COVID-19).¹ Personal protective equipment (PPE), including N95 respirators (N95s), is essential for prevention of COVID-19. The Centers for Disease Control and Prevention recommends that health care workers dispose of N95s after a single patient encounter. However, it recommends N95 extended use (wearing the same N95 for multiple patient encounters) and limited reuse (storing an N95 between encounters for use over multiple encounters) during critical PPE shortages.^{2,3} There are limited data regarding N95 reuse and extended use. Existing studies were conducted in laboratories, not clinical environments.^{4,5} Inadequate supplies of N95s have forced many emergency departments to implement various N95 reuse and extended use policies but without empirical evidence of their effectiveness. We examined the prevalence of N95 fit test failure while reusing 2 common types of N95 masks.

Methods | We performed a cross-sectional study of N95 fit at the University of California, San Francisco (UCSF) emergency department from April 4 to April 6, 2020. We enrolled a convenience sample of health care workers (physicians, nurses, nurse practitioners, physician assistants, and patient care technicians) on their clinical shifts when the researchers



Audio